

Wall crossing for symplectic vortices and quantum cohomology

Kai Cieliebak*

Ludwig-Maximilians-Universität München

Dietmar A. Salamon

ETH-Zürich

18 September 2002

Abstract

We derive a wall crossing formula for the symplectic vortex invariants of toric manifolds. As an application, we give a proof of Batyrev's formula for the quantum cohomology of a monotone toric manifold with minimal Chern number at least two.

1 Introduction

Let T be a torus of dimension k , denote by $\mathfrak{t}$ its Lie algebra, by

$$\Lambda := \{\xi \in \mathfrak{t} \mid \exp(\xi) = 1\}$$

the integer lattice, and by

$$\Lambda^* := \{w \in \mathfrak{t}^* \mid \langle w, \xi \rangle \in \mathbb{Z} \text{ for } \xi \in \Lambda\}$$

the dual lattice. Suppose T acts diagonally on $\mathbb{C}^n$. The action is determined by n homomorphisms $\rho_\nu : T \rightarrow S^1$, $\nu = 1, \dots, n$. We write each homomorphism ρ_ν in the form

$$\rho_\nu(\exp(\xi)) = e^{-2\pi i \langle w_\nu, \xi \rangle}, \quad w_\nu \in \Lambda^*.$$

The moment map of this action, with respect to the standard symplectic form on $\mathbb{C}^n$, is given by

$$\mu(x) = \pi \sum_{\nu=1}^n |x_\nu|^2 w_\nu \tag{1}$$

*Supported by National Science Foundation Grant DMS-0072267

for $x = (x_1, \dots, x_n) \in \mathbb{C}^n$. We assume throughout that μ is proper and that the vectors w_ν span the space $\mathfrak{t}^*$. In [5] we have defined invariants

$$\Phi_\lambda^{\rho, \tau} : S^m(\mathfrak{t}^*) \rightarrow \mathbb{R}, \quad m := n - \dim T + \sum_{\nu=1}^n d_\nu \geq 0, \quad (2)$$

by counting solutions of the genus zero symplectic vortex equations (see Section 2). Here $\lambda \in \Lambda$, $d_\nu := \langle w_\nu, \lambda \rangle$, τ is a regular value of the moment map, and $S^m(\mathfrak{t}^*)$ denotes the space of real valued polynomials of degree m on $\mathfrak{t}$. Note that $S^*(\mathfrak{t}^*)$ is canonically isomorphic to the cohomology $H^*(BT; \mathbb{R})$ of the classifying space $BT = ET/T$. The isomorphism takes $w_\nu \in \mathfrak{t}^*$ to the first Chern class of the bundle $ET \times_{\rho_\nu} \mathbb{C} \rightarrow BT$. The invariant $\Phi_\lambda^{\rho, \tau}$ takes rational values on integral cohomology classes. These correspond to polynomials that map the lattice to the integers.

An element $\tau \in \mathfrak{t}^*$ is a singular value of μ if and only if it can be expressed as a positive linear combination of at most $k-1$ of the vectors w_ν . The set of singular values is a disjoint union of open cones of codimensions 1 to k . A cone of codimension j is called a **wall of codimension j** .

Let $\tau_0 \in \mathfrak{t}^*$ be an element of a wall of codimension one, $\tau_1 \in \mathfrak{t}^*$ be transverse to the wall at τ_0 , and $e_1 \in \Lambda$ be the unique primitive lattice vector that is orthogonal to the wall at τ_0 and satisfies $\langle \tau_1, e_1 \rangle > 0$. Denote by $T_1 \subset T$ the subtorus generated by e_1 and by $\mathfrak{t}_1$ its Lie algebra. Let

$$I := \{\nu \mid \langle w_\nu, e_1 \rangle = 0\}.$$

The action ρ induces an action ρ_0 of the quotient torus

$$T_0 := T/T_1$$

on the space

$$\mathbb{C}^I := \{x \in \mathbb{C}^n \mid x_\nu = 0 \text{ for } \nu \notin I\}.$$

The moment map of this action is the restriction $\mu_0 := \mu|_{\mathbb{C}^I} : \mathbb{C}^I \rightarrow \mathfrak{t}_0^* := \mathfrak{t}_1^\perp$. The following *wall crossing formula* expresses the difference of the invariants on the two sides of the wall as the invariant of the reduced problem at τ_0 .

Theorem 1.1 (Genus Zero Wall Crossing). *Let $\alpha \in S^*(\mathfrak{t}^*)$, $\lambda \in \Lambda$, and $d_\nu := \langle w_\nu, \lambda \rangle$. Then for every sufficiently small positive number ε we have*

$$\Phi_\lambda^{\rho, \tau_0 + \varepsilon \tau_1}(\alpha) - \Phi_\lambda^{\rho, \tau_0 - \varepsilon \tau_1}(\alpha) = \Phi_{\lambda_0}^{\rho_0, \tau_0}(\alpha_0),$$

where λ_0 is the projection of λ to $\mathfrak{t}_0 := \mathfrak{t}/\mathfrak{t}_1$ and

$$\alpha_0(\xi) = \frac{1}{2\pi i} \oint \frac{\alpha(\xi + ze_1)}{\prod_{\nu \notin I} \langle w_\nu, \xi + ze_1 \rangle^{d_\nu + 1}} dz.$$

Here for each ξ the integral is understood over a circle in the complex plane enclosing all the poles of the integrand.

There is an analogous wall crossing formula for higher genus which is formulated in Theorem 3.1 below.

Theorem 1.1 gives rise to an explicit formula for the genus zero invariants. To formulate the result we introduce the following notation. For a tuple of nonnegative integers $\ell = (\ell_1, \dots, \ell_n)$ denote

$$\mathbf{w}^\ell := \mathbf{w}_1^{\ell_1} \cdots \mathbf{w}_n^{\ell_n} \in S^{|\ell|}(\mathfrak{t}^*), \quad |\ell| := \ell_1 + \cdots + \ell_n.$$

Given such a tuple ℓ and a lattice vector λ we introduce the set $\mathcal{I}_\lambda(\ell)$ of partitions $\{1, \dots, n\} = I_1 \cup \cdots \cup I_k$ that satisfy the following two conditions.

(Dimension) For every $j \in \{1, \dots, k\}$, the subspace

$$E_j := \text{span}\{\mathbf{w}_\nu \mid \nu \in I_1 \cup \cdots \cup I_j\} \subset \mathfrak{t}^*$$

has dimension j and $\mathbf{w}_\nu \notin E_j$ for every $\nu \in I_{j+1} \cup \cdots \cup I_k$.

(Degree) For every $j \in \{1, \dots, k\}$,

$$\sum_{\nu \in I_j} (\ell_\nu - d_\nu - 1) = -1. \quad d_\nu := \langle \mathbf{w}_\nu, \lambda \rangle.$$

Theorem 1.2 (Genus Zero Invariants). *Let $\lambda \in \Lambda$, $d_\nu := \langle \mathbf{w}_\nu, \lambda \rangle$, and ℓ be an n -tuple of nonnegative integers.*

(i) *If $\mathcal{I}_\lambda(\ell) = \emptyset$ then $\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = 0$ for all τ .*

(ii) *Let $J \subset \{1, \dots, n\}$ be a subset with k elements such that $\{\mathbf{w}_\nu \mid \nu \in J\}$ is a basis of $\mathfrak{t}^*$ and assume*

$$\ell_\nu = \begin{cases} d_\nu & \text{if } \nu \in J, \\ d_\nu + 1 & \text{if } \nu \notin J. \end{cases}$$

If τ belongs to the cone $C(J)$ spanned by $\{\mathbf{w}_\nu \mid \nu \in J\}$ then

$$\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = \frac{1}{|\det(\langle \mathbf{w}_\nu, e_j \rangle_{\nu \in J, j=1, \dots, k})|},$$

where $e_1, \dots, e_k$ is any basis of the lattice Λ . Otherwise $\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = 0$.

(iii) *Let $\lambda' \in \Lambda$ and define $d'_\nu := \langle \mathbf{w}_\nu, \lambda' \rangle$. If $\ell_\nu + d'_\nu \geq 0$ for every ν then*

$$\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = \Phi_{\lambda+\lambda'}^{\rho, \tau}(\mathbf{w}^{\ell+d'}).$$

(iv) *Assume $d_\nu \geq -1$ for every ν . Then every element of $S^*(\mathfrak{t}^*)$ is a linear combination of monomials $\mathbf{w}^\ell$ that satisfy either (i) or (ii).*

(v) *Let $J_\ell := \{\nu \mid \ell_\nu \leq d_\nu\}$. If $\tau \notin C(J_\ell)$ then $\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = 0$.*

Remark. Assertions (i), (ii) and (iv) can be used to compute the genus zero invariants whenever $d_\nu \geq -1$. This restriction can be removed by using (iii).

Now assume that T acts freely on $\mu^{-1}(\tau)$, where

$$\tau := \sum_{\nu=1}^n w_\nu,$$

and that the symplectic quotient

$$\bar{M} := \mathbb{C}^n // T(\tau) := \mu^{-1}(\tau)/T$$

has minimal Chern number $N := \max \{m \in \mathbb{Z} \mid \tau/m \in \Lambda^*\} \geq 2$. These conditions guarantee that the symplectic quotient $\bar{M}$ is a monotone toric manifold. Combining Theorem 1.2 with the results of [8] one can compute the genus zero Gromov–Witten invariants of the symplectic quotient. More precisely, denote by $S^*(\mathfrak{t}^*) \rightarrow H^*(\bar{M}; \mathbb{R}) : \alpha \mapsto \bar{\alpha}$ the Kirwan homomorphism. Consider the dual homomorphism $H_2(\bar{M}; \mathbb{Z}) \rightarrow \Lambda$ in degree two. This homomorphism is injective. We denote its image by $\Lambda(\tau)$ and the inverse map by $\Lambda(\tau) \rightarrow H_2(\bar{M}; \mathbb{Z}) : \lambda \mapsto \bar{\lambda}$. Given $\lambda \in \Lambda(\tau)$ we denote by $\text{GW}_{\bar{\lambda}}^{\bar{M}}$ the genus zero Gromov–Witten invariant of $\bar{M}$ with fixed marked points in the homology class $\bar{\lambda}$. In [8, Theorem A] it is proved that, for every $\lambda \in \Lambda(\tau)$ and every n -tuple $\ell = (\ell_1, \dots, \ell_n)$ of nonnegative integers,

$$\Phi_{\lambda}^{\rho, \tau}(w^\ell) = \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}_1, \dots, \bar{w}_1, \dots, \bar{w}_n, \dots, \bar{w}_n), \quad (3)$$

where each argument $\bar{w}_\nu$ occurs ℓ_ν times. Thus Theorem 1.2 allows us to compute the genus zero Gromov–Witten invariants of tuples of cohomology classes of degree two. This can be used to compute the quantum cohomology ring of the symplectic quotient. The statement of the theorem requires some preparation.

The **chamber** $C(\tau)$ is defined as the component of the set of regular values of μ that contains τ . The **effective cone** $\Lambda_{\text{eff}}(\tau) \subset \Lambda(\tau)$ is defined as the set of lattice vectors $\lambda \in \Lambda(\tau)$ that satisfy $\langle \tau', \lambda \rangle \geq 0$ for every $\tau' \in C(\tau)$.

Let $\mathcal{R}$ be any graded commutative algebra (over the reals) with unit which is equipped with a homomorphism

$$\Lambda_{\text{eff}}(\tau) \rightarrow \mathcal{R} : \lambda \mapsto q^\lambda$$

from the additive semigroup $\Lambda_{\text{eff}}(\tau)$ to the multiplicative semigroup $\mathcal{R}$ such that $\deg(q^\lambda) = 2\langle \tau, \lambda \rangle$. Given such a graded algebra define the quantum cohomology ring $\text{QH}^*(\bar{M}; \mathcal{R})$ as the tensor product

$$\text{QH}^*(\bar{M}; \mathcal{R}) := H^*(\bar{M}; \mathbb{R}) \otimes \mathcal{R}$$

(of vector spaces over the reals). Thus an element of $\text{QH}^*(\bar{M}; \mathcal{R})$ is a finite sum $\bar{\alpha} = \sum_{r \in \mathcal{R}} \bar{\alpha}_r r$ where $\bar{\alpha}_r \in H^*(\bar{M}; \mathbb{R})$. The ring structure is defined by

$$\bar{\alpha}' * \bar{\alpha}'' := \sum_i \sum_{\lambda \in \Lambda_{\text{eff}}(\tau)} \sum_{r', r''} \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{\alpha}'_{r'}, \bar{\alpha}''_{r''}, \bar{e}_i) \bar{e}_i^* r' r'' q^\lambda,$$

where the $\bar{e}_i$ form a basis of $H^*(\bar{M}; \mathbb{R})$ and the $\bar{e}_i^*$ denote the dual basis with respect to the cup product pairing.

Theorem 1.3 (Quantum Cohomology). *Let $\bar{M} = \mathbb{C}^n // T(\tau)$ be a (nonempty) monotone toric manifold with minimal Chern number $N \geq 2$. Then the ring homomorphism*

$$\mathcal{R}[u_1, \dots, u_n] \rightarrow \mathrm{QH}^*(\bar{M}; \mathcal{R}) : ru^\ell \mapsto r\bar{w}^{*\ell} \quad (4)$$

induces an isomorphism $\mathrm{QH}^*(\bar{M}; \mathcal{R}) \cong \mathcal{R}[u_1, \dots, u_n]/\mathcal{J}$, where the ideal $\mathcal{J} \subset \mathcal{R}[u_1, \dots, u_n]$ is generated by the relations

$$\begin{aligned} \sum_{\nu=1}^n \eta_\nu w_\nu = 0 &\implies \sum_{\nu=1}^n \eta_\nu u_\nu = 0, \\ \tau \notin C(\{1, \dots, n\} \setminus \{\nu\}) &\implies u_\nu = 0, \\ \lambda \in \Lambda_{\mathrm{eff}}(\tau), \quad d_\nu^\pm := \max\{\pm \langle w_\nu, \lambda \rangle, 0\} &\implies \prod_\nu u_\nu^{d_\nu^+} = q^\lambda \prod_\nu u_\nu^{d_\nu^-}. \end{aligned}$$

The ring $\mathcal{R}[u_1, \dots, u_n]/\mathcal{J}$ was introduced by Batyrev [4]. It also appeared in Givental's work on mirror symmetry [9] for the monotone case. Examples of Spielberg [13, 14] show that, in the nonmonotone case, the kernel of the homomorphism (4) is not necessarily equal to $\mathcal{J}$. For special cases the isomorphism $\mathcal{R}[u_1, \dots, u_n]/\mathcal{J} \rightarrow \mathrm{QH}^*(\bar{M}; \mathcal{R})$ was established in [12, 13]. The reason for our hypothesis $N \geq 2$ lies in the identity (3) which, in general, does not continue to hold in the case $N = 1$ (the degrees of all the classes must be less than twice the minimal Chern number).

In Section 2 we explain some background from [5] about the symplectic vortex equations. The wall crossing formula (for arbitrary genus) is restated in Section 3 and proved in Section 4. We prove Theorem 1.2 in Section 5 and Theorem 1.3 in Section 6.

2 The symplectic vortex equations

Fix a compact Riemann surface $(\Sigma, j_\Sigma, \mathrm{dvol}_\Sigma)$, a principal T -bundle $P \rightarrow \Sigma$, and an invariant inner product on $\mathfrak{t}$. The characteristic vector of P will be denoted by

$$\lambda(P) := \int_\Sigma F_A \in \Lambda.$$

Here $A \in \mathcal{A} = \mathcal{A}(P)$ is a connection on P and $\lambda(P)$ is independent of the choice of the connection. The symplectic vortex equations (at a parameter $\tau \in \mathfrak{t}^*$) have the form

$$\bar{\partial}_A u_\nu = 0, \quad *F_A + \pi \sum_{\nu=1}^n |u_\nu|^2 w_\nu = \frac{*_{\mathfrak{t}} \lambda(P)}{\mathrm{Vol}(\Sigma)} + \tau, \quad (5)$$

where $u_\nu : P \rightarrow \mathbb{C}$ is equivariant with respect to ρ_ν , $* : \Omega^2(\Sigma, \mathfrak{t}) \rightarrow \Omega^0(\Sigma, \mathfrak{t}^*)$ denotes the Hodge $*$ -operator determined by the volume form on Σ and the inner product on $\mathfrak{t}$, and $*_{\mathfrak{t}} : \mathfrak{t} \rightarrow \mathfrak{t}^*$ denotes the isomorphism induced by the

inner product. The gauge group $\mathcal{G} := C^\infty(\Sigma, T)$ acts on the space of solutions of (5) with finite isotropy if and only if τ is a regular value of μ . Moreover, the moduli space

$$\mathcal{M}(\tau) := \{(A, u) \mid u \text{ and } A \text{ satisfy (5)}\} / \mathcal{G}_0$$

of based gauge equivalence classes of solutions of (5) is compact (see [5]). Here we fix a point $z_0 \in \Sigma$ and denote the based gauge group by

$$\mathcal{G}_0 := \{g \in \mathcal{G} \mid g(z_0) = \mathbb{1}\}.$$

Think of this moduli space as a subset of the space

$$\mathcal{B} := \frac{\mathcal{A}(P) \times C_T^\infty(P, \mathbb{C}^n)}{\mathcal{G}_0},$$

where $g \in \mathcal{G}_0$ acts by $g^*(A, u) := (A + g^{-1}dg, \rho(g)^{-1}u)$. The group T (of constant gauge transformations) acts contravariantly on $\mathcal{B}$ and we identify

$$\mathcal{B} \times_T ET \cong \frac{\mathcal{A}(P) \times C_T^\infty(P, \mathbb{C}^n) \times ET}{\mathcal{G}}$$

where $\mathcal{G}$ acts by $g^*(A, u, e) := (A + g^{-1}dg, \rho(g)^{-1}u, g(z_0)^{-1}e)$. The invariants introduced in [5] are obtained by integrating equivariant cohomology classes of $\mathcal{B}$ over $\mathcal{M}(\tau)/T$. Here integration is to be understood in terms of evaluating the equivariant Euler class of an associated T -moduli problem on the equivariant cohomology class in question (see [6]).

It is useful to think of u_ν as a section of the complex line bundle

$$L_\nu := P \times_{\rho_\nu} \mathbb{C} \rightarrow \Sigma, \quad (6)$$

where the equivalence relation on $P \times \mathbb{C}$ is $[p, \zeta] \equiv [pg, \rho_\nu(g)^{-1}\zeta]$ for $g \in T$. This bundle has degree

$$d_\nu := \frac{i}{2\pi} \dot{\rho}_\nu(\lambda(P)) = \langle w_\nu, \lambda(P) \rangle$$

and the moduli space $\mathcal{M}(\tau)/T$ has virtual dimension

$$\dim_{\mathbb{R}} \mathcal{M}(\tau)/T = (n - \dim T)(2 - 2g) + 2 \sum_{\nu=1}^n d_\nu =: 2m,$$

where g is the genus of Σ . Integration over the moduli space gives rise to a homomorphism

$$\Phi_{\lambda, g}^{\rho, \tau} : S^*(\mathfrak{t}^*) \otimes H^*(\mathcal{A}/\mathcal{G}_0) \rightarrow \mathbb{R}.$$

Formally, this homomorphism can be written in the form

$$\Phi_{\lambda, g}^{\rho, \tau}(\alpha) := \int_{\mathcal{M}(\tau)/T} \pi^* \alpha \quad (7)$$

for $\alpha \in S^*(\mathfrak{t}^*) \otimes H^*(\mathcal{A}/\mathcal{G}_0) \cong H_T^*(\mathcal{A}/\mathcal{G}_0)$. Here $\pi : \mathcal{B} \rightarrow \mathcal{A}/\mathcal{G}_0$ is the obvious projection and $\pi^* : H_T^*(\mathcal{A}/\mathcal{G}_0) \rightarrow H_T^*(\mathcal{B})$ is the induced homomorphism on equivariant cohomology. The precise definition of the invariant involves the equivariant

Euler class of the T -moduli problem associated to equation (5) (see [6, 5]). The invariant $\Phi_{\lambda, g}^{\rho, \tau}(\alpha)$ can only be nonzero when α has degree $2m$ and it takes rational values on cohomology classes that take the integer lattice in $\mathfrak{t}$ to the integral cohomology of $\mathcal{A}/\mathcal{G}_0$.

3 Wall crossing

In this section we formulate the wall crossing formula for arbitrary genus at an element $\tau_0 \in \mathfrak{t}^*$ of a wall of codimension one. This means that there exists an index set $I \subset \{1, \dots, n\}$ satisfying the following conditions.

- (i) The subspace $W_I := \text{span}\{\mathbf{w}_\nu \mid \nu \in I\} \subset \mathfrak{t}^*$ has dimension $k-1$ and $\mathbf{w}_\nu \notin W_I$ for every $\nu \notin I$.
- (ii) $\tau_0 \in \mu(\mathbb{C}^I)$, where $\mathbb{C}^I := \{x \in \mathbb{C}^n \mid x_\nu = 0 \text{ for } \nu \notin I\}$.
- (iii) If $J \subset \{1, \dots, n\}$ is another index set satisfying (i) then $\tau_0 \notin \mu(\mathbb{C}^J)$.

Note that under these conditions τ_0 is a positive linear combination of precisely $k-1$ linearly independent vectors from the set $\{\mathbf{w}_\nu \mid \nu \in I\}$. Choose a vector $\tau_1 \in \mathfrak{t}^*$ that is transverse to $\mu(\mathbb{C}^I)$ and let $e_1 \in \Lambda$ be the unique primitive lattice vector that satisfies $\langle \tau_1, e_1 \rangle > 0$ and is orthogonal to the wall at τ_0 :

$$\langle \mathbf{w}_\nu, e_1 \rangle = 0 \quad \text{for } \nu \in I.$$

Denote by $T_1 \subset T$ the subtorus generated by e_1 and by $\mathfrak{t}_1$ its Lie algebra. Let $T_0 := T/T_1$ be the quotient torus and $\mathfrak{t}_0 := \mathfrak{t}/\mathfrak{t}_1$ be its Lie algebra. Then the action ρ induces an action ρ_0 of T_0 on $\mathbb{C}^I$.

The wall crossing number will be expressed as an integral over the moduli space $\mathcal{M}_0$ of based gauge equivalence classes of solutions $(A, \{u_\nu\}_{\nu \in I})$ of the equations

$$\bar{\partial}_A u_\nu = 0 \quad (\nu \in I), \quad *F_A + \pi \sum_{\nu \in I} |u_\nu|^2 \mathbf{w}_\nu = \frac{*_{\mathfrak{t}} \lambda(P)}{\text{Vol}(\Sigma)} + \tau_0. \quad (8)$$

We shall view this as a T_0 -moduli problem. Indeed, the subgroup $T_1 \subset T$ acts trivially on $\mathcal{M}_0$. However, since τ_0 is a regular value of $\mu|_{\mathbb{C}^I}$, the quotient group $T_0 = T/T_1$ acts on $\mathcal{M}_0$ with finite isotropy.

It is interesting to compare $\mathcal{M}_0$ with the moduli space $\mathcal{M}(\mathbb{C}^I, P_0, \tau_0)$ of based gauge equivalence classes of solutions of (5) with $\mathbb{C}^n$, T , and P replaced $\mathbb{C}^I$, $T_0 := T/T_1$, and $P_0 := P/T_1$. There is a natural projection

$$\mathcal{M}_0 \longrightarrow \mathcal{M}(\mathbb{C}^I, P_0, \tau_0) : [A, \{u_\nu\}_{\nu \in I}] \mapsto [A_0, \{u_\nu\}_{\nu \in I}], \quad (9)$$

where $A_0 := \Pi_0 A \in \Omega^1(P, \mathfrak{t}_0)$ can be thought of as a connection on P_0 . Here $\Pi_0 : \mathfrak{t} \rightarrow \mathfrak{t}_0$ denotes the canonical projection. If $(A, \{u_\nu\}_{\nu \in I})$ satisfies (8) then the tuple $(A_0, \{u_\nu\}_{\nu \in I})$ satisfies the equations

$$\bar{\partial}_{A_0} u_\nu = 0 \quad (\nu \in I), \quad *_0 F_{A_0} + \pi \sum_{\nu \in I} |u_\nu|^2 \mathbf{w}_\nu = \frac{*_{\mathfrak{t}_0} \lambda(P_0)}{\text{Vol}(\Sigma)} + \tau_0.$$

and hence belongs to the moduli space $\mathcal{M}(\mathbb{C}^I, P_0, \tau_0)$. Here $*_{t_0} : \mathfrak{t}/\mathfrak{t}_1 \rightarrow \mathfrak{t}_1^\perp$ is given by $[\xi] \mapsto \xi - |e_1|^{-2} \langle \xi, e_1 \rangle e_1$ and $*_0 : \Omega^2(\Sigma, \mathfrak{t}/\mathfrak{t}_1) \rightarrow \Omega^0(\Sigma, \mathfrak{t}_1^\perp)$ is induced by the Hodge $*$ -operator on Σ and $*_{t_0}$. The map (9) defines a fibration whose fiber can be described as follows.

Choose a complement of $\mathfrak{t}_1$ in $\mathfrak{t}$ and denote the resulting projection by $\Pi_1 : \mathfrak{t} \rightarrow \mathfrak{t}_1$. Define Pic_1 as the space of based gauge equivalence classes of real valued T -invariant 1-forms $A_1 \in \Omega^1(P)$ that satisfy $A_1(pe_1) = 1$ for every $p \in P$ and

$$*F_{A_1} = \frac{\Pi_1 \lambda(P)}{\text{Vol}(\Sigma)}. \quad (10)$$

Here the based gauge group is $\mathcal{G}_1 := \{g_1 : \Sigma \rightarrow T_1 \mid g_1(z_0) = \mathbb{1}\}$ and it acts by $(g_1^* A_1)e_1 := A_1 e_1 + g_1^{-1} dg_1$. Note that Pic_1 is a $2g$ -torus. It is the fibre in (9) because the subgroup $\mathcal{G}_1 \subset \mathcal{G}_0$ acts trivially on $A_0 = \Pi_0 A$ and u_ν for every $\nu \in I$. We emphasize that the fibration (9) need not be a product. The reason is that the map $A \mapsto A_1 := \Pi_1 A$ will not, in general, be gauge invariant. We are now in a position to state the wall crossing formula for arbitrary genus. Think of $\mathcal{M}_0$ as a subset of the space $\mathcal{B}_0 := \mathcal{A}(P) \times C_T^\infty(P, \mathbb{C}^I)/\mathcal{G}_0$ and denote by $\pi_0^* : S^*(\mathfrak{t}_0^*) \otimes H^*(\mathcal{A}/\mathcal{G}_0) \rightarrow H_{T_0}^*(\mathcal{B}_0)$ the homomorphism on equivariant cohomology induced by the T_0 -invariant projection $\pi_0 : \mathcal{B}_0 \rightarrow \mathcal{A}/\mathcal{G}_0$.

Theorem 3.1 (Wall Crossing). *Let $\alpha \in S^*(\mathfrak{t}^*) \otimes H^*(\mathcal{A}/\mathcal{G}_0)$, $\lambda \in \Lambda$, and $d_\nu := \langle w_\nu, \lambda \rangle$. Then, for every sufficiently small positive number ε , we have*

$$\Phi_{\lambda, g}^{\rho, \tau_0 + \varepsilon \tau_1}(\alpha) - \Phi_{\lambda, g}^{\rho, \tau_0 - \varepsilon \tau_1}(\alpha) = \int_{\mathcal{M}_0/T_0} \pi_0^* \alpha_0, \quad (11)$$

where $\alpha_0 \in S^*(\mathfrak{t}_0^*) \otimes H^*(\mathcal{A}/\mathcal{G}_0)$ is the polynomial map defined by

$$\alpha_0(\xi) := \frac{1}{2\pi i} \oint \frac{\alpha(\xi + ze_1)}{\prod_{\nu \notin I} \langle w_\nu, \xi + ze_1 \rangle^{d_\nu + 1 - g}} \exp \left(\sum_{\nu \notin I} \frac{\Omega_\nu}{\langle w_\nu, \xi + ze_1 \rangle} \right) dz.$$

Here Ω_ν is the closed 2-form on $\mathcal{A}/\mathcal{G}_0$ defined by

$$\Omega_\nu := \sum_{j, j'=1}^k \Omega_{jj'} w_{\nu j} w_{\nu j'}.$$

Here we have chosen a basis $e_1, \dots, e_k$ of $\mathfrak{t}$ and a symplectic basis $\alpha_1, \dots, \alpha_{2g}$ of $H^1(\Sigma; \mathbb{Z})$. These bases induce a basis τ_{ij} of $H^1(\mathcal{A}/\mathcal{G}_0; \mathbb{Z})$ and $\Omega_{jj'}$ and $w_{\nu j}$ are defined by

$$\Omega_{jj'} := \sum_{i=1}^g \tau_{ij} \wedge \tau_{i+g, j'}, \quad w_{\nu j} := \langle w_\nu, e_j \rangle.$$

Note that $\Omega_{jj'}$ is independent of the choice of the α_i and Ω_ν is independent of the choice of both bases. For each ξ , the integral in the definition of $\alpha_0(\xi)$ is over a circle in the complex plane enclosing all the poles of the integrand. The integral in (11) is understood as the evaluation of the equivariant Euler class of the T_0 -moduli problem associated to (8) on the class $\pi_0^* \alpha_0$ (see [6]).

Remark 3.2 (Residues). Consider a rational function $f : \mathbb{C} \rightarrow \mathbb{C}$ with poles $p_1, \dots, p_n$. It induces a meromorphic 1-form $f dz$. Let $\oint f dz$ be the integral of $f dz$ over a closed curve in $\mathbb{C}$ around all the poles of f . By the residue theorem,

$$\frac{1}{2\pi i} \oint f dz = -\text{Res}_\infty(f dz) = \sum_{j=1}^n \text{Res}_{p_j}(f dz).$$

Note that the 1-form $f dz$ and hence the residue at infinity do not change if the complex coordinate is shifted by $z \mapsto z + c$. If we expand f as a Laurent series

$$f(z) = \sum_{k=-\infty}^{k_0} a_k z^k$$

in z^{-1} that converges near infinity then the residue at infinity is minus the coefficient of z^{-1} , i.e. $\text{Res}_\infty(f) = -a_{-1}$.

Example 3.3. Consider the action of the 1-torus $T = \mathbb{R}/\mathbb{Z}$ on $\mathbb{C}^n$ with positive integer weights $w_\nu = \ell_\nu \in \Lambda^* \cong \mathbb{Z}$. Let $c \in \Lambda^*$ be the standard generator $c(\xi) = \xi$ and pick a homology class $\lambda = d \in \Lambda \cong \mathbb{Z}$. Assume

$$m := \sum_{\nu=1}^n (d\ell_\nu + 1 - g) + g - 1 \geq 0.$$

We compute the invariant $\Phi_{d,g}^{\rho,\tau}(c^m)$ in the nonempty chamber by wall crossing from the empty chamber. Here $I = \emptyset$, $T_1 = T = \mathbb{R}/\mathbb{Z}$, $e_1 = 1$, and $T_0 = \{\mathbb{1}\}$. Then $\mathcal{M}_0 = \text{Pic}$ is a $2g$ -torus and $\Omega_\nu = \ell_\nu^2 \Omega$, where Ω is the standard symplectic form on Pic . It satisfies

$$\frac{1}{g!} \int_{\text{Pic}} \Omega^g = 1.$$

The integrand in Theorem 3.1 is given by

$$\alpha_0 = \frac{1}{2\pi i} \oint \frac{z^m}{\prod_{\nu} (\ell_\nu z)^{d\ell_\nu + 1 - g}} \exp\left(\frac{(\ell_1 + \dots + \ell_n)\Omega}{z}\right) = \frac{(\ell_1 + \dots + \ell_n)^g \Omega^g}{\prod_{\nu=1}^n \ell_\nu^{d\ell_\nu + 1 - g} g!}.$$

Integrating this class over Pic yields the formula from [5]:

$$\Phi_{\lambda,g}^{\rho,\tau}(c^m) = \frac{(\ell_1 + \dots + \ell_n)^g}{\prod_{\nu=1}^n \ell_\nu^{d\ell_\nu + 1 - g}}.$$

4 Proof of the wall crossing formula

4.1 A cobordism argument

We introduce the gauge invariant differential equations

$$\begin{aligned} \bar{\partial}_A u_\nu &= 0 \quad (\nu = 1, \dots, n), \\ *F_A + \pi \sum_{\nu \in I} |u_\nu|^2 w_\nu &= \frac{*_{\mathfrak{t}} \lambda(P)}{\text{Vol}(\Sigma)} + \tau_0, \quad \sum_{\nu \notin I} \|u_\nu\|^2 = 1. \end{aligned} \tag{12}$$

Denote the moduli space of based gauge equivalence classes of solutions of (12) by

$$\mathcal{S}_0 := \{(A, u) \mid A \text{ and } u \text{ satisfy (12)}\} / \mathcal{G}_0.$$

Note that there is a T -equivariant projection $\mathcal{S}_0 \rightarrow \mathcal{M}_0$ whose fiber is the unit sphere in the kernel of the Cauchy–Riemann operator in the variables u_ν for $\nu \notin I$.

Proposition 4.1. *The wall crossing number can be expressed in the form*

$$\Phi_{\lambda, g}^{\rho, \tau_0 + \varepsilon \tau_1}(\alpha) - \Phi_{\lambda, g}^{\rho, \tau_0 - \varepsilon \tau_1}(\alpha) = - \int_{\mathcal{S}_0/T} \pi^* \alpha,$$

where the orientation of $\mathcal{S}_0/T$ is determined by e_1 as in Remark 4.2.

Remark 4.2. The moduli space $\mathcal{S}_0/T$ is oriented as follows. Denote

$$\mathcal{X} := \Omega^1(\Sigma, \mathfrak{t}) \oplus \bigoplus_{\nu=1}^n \Omega^0(\Sigma, L_\nu), \quad \mathcal{Y} := \Omega^0(\Sigma, \mathfrak{t}^* \otimes \mathbb{C}) \oplus \bigoplus_{\nu=1}^n \Omega^{0,1}(\Sigma, L_\nu),$$

where $L_\nu \rightarrow \Sigma$ is the line bundle (6). Define the operator $\mathcal{D}_0 : \mathcal{X} \rightarrow \mathcal{Y}$ by

$$\mathcal{D}_0(\alpha, \hat{u}) := \begin{pmatrix} *d\alpha + 2\pi \sum_{\nu \in I} \langle u_\nu, \hat{u}_\nu \rangle w_\nu \\ d^* \alpha + 2\pi \sum_{\nu \in I} \langle iu_\nu, \hat{u}_\nu \rangle w_\nu \\ \bar{\partial}_A \hat{u}_\nu + \rho_\nu(\alpha)^{0,1} u_\nu \end{pmatrix}.$$

The first component is the real part in $\Omega^0(\Sigma, \mathfrak{t}^* \otimes \mathbb{C})$ (it is the linearization of the second equation in (12)) and the second component is the imaginary part (it corresponds to the local slice condition for action of $\mathcal{G}/T_1$). The third component is the linearization of the first equation in (12). The operator $\mathcal{D}_0$ is complex linear, where the complex structure on $\Omega^1(\Sigma, \mathfrak{t})$ is given by the Hodge $*$ -operator $\alpha \mapsto *\alpha = -\alpha \circ J_\Sigma$. Define the linear functional $\Psi : \mathcal{X} \rightarrow \mathbb{C}$ by

$$\Psi(\alpha, \hat{u}) := \frac{2\pi}{\langle \tau_1, e_1 \rangle \text{Vol}(\Sigma)} \sum_{\nu \notin I} \langle w_\nu, e_1 \rangle \int_\Sigma (\langle u_\nu, \hat{u}_\nu \rangle + i \langle iu_\nu, \hat{u}_\nu \rangle) \text{dvol}_\Sigma.$$

The imaginary part of Ψ corresponds to the local slice for the T_1 -action.

Note that the cokernel of $\mathcal{D}_0$ always contains the constant functions in $\Omega^0(\Sigma, \mathfrak{t}_1 \otimes \mathbb{C})$. We assume for simplicity that the cokernel is equal to this space and that

$$\Psi(0, \{u_\nu\}_{\nu \notin I}) = \frac{2\pi}{\langle \tau_1, e_1 \rangle \text{Vol}(\Sigma)} \sum_{\nu \notin I} \|u_\nu\|_{L^2}^2 \langle w_\nu, e_1 \rangle \neq 0.$$

Then $\mathcal{S}_0$ is a smooth manifold near $[A, u]$ and the tangent space of $\mathcal{S}_0/T$ is

$$T_{[A, u]} \mathcal{S}_0/T = \left\{ (\alpha, \hat{u}) \mid \mathcal{D}_0(\alpha, \hat{u}) = 0, \sum_{\nu \notin I} \langle u_\nu, \hat{u}_\nu \rangle = 0, \text{Im } \Psi(\alpha, \hat{u}) = 0 \right\}.$$

A basis $v_1, \dots, v_{2m}$ of the tangent space is called **positively oriented** if the vectors $v_1, \dots, v_{2m}$, $w_0 := (0, \{u_\nu\}_{\nu \notin I})$, $w_1 := (0, \{-2\pi i \langle w_\nu, e_1 \rangle u_\nu\}_{\nu \notin I})$ form a positive basis of the complex vector space $\ker \mathcal{D}_0$. Note that the orientation depends on e_1 and that $m = (n - \dim T)(1 - g) + \sum_{\nu=1}^n d_\nu$.

Proof of Proposition 4.1. Denote $\tau_t := \tau_0 + t\tau_1$ and consider the moduli space

$$\mathcal{W} := \{(t, A, u) \mid -\varepsilon \leq t \leq \varepsilon, (5) \text{ holds with } \tau = \tau_t\} / \mathcal{G}_0.$$

This space has boundary

$$\partial \mathcal{W} = (\{-\varepsilon\} \times \mathcal{M}(\tau_{-\varepsilon})) \cup (\{\varepsilon\} \times \mathcal{M}(\tau_\varepsilon)).$$

Now the group T of constant gauge transformations does not act with finite isotropy on $\mathcal{W}$. Call an element of $\mathcal{W}$ **singular** if its isotropy subgroup has positive dimension. Then the singular subset of $\mathcal{W}$ is precisely the moduli space $\mathcal{M}_0$ introduced in the previous section. Cutting out a neighbourhood of this singular set we obtain the subset $\mathcal{W}_\delta \subset \mathcal{W}$ of all (based gauge equivalence classes of) triples $[t, A, u] \in \mathcal{W}$ that satisfy $\sum_{\nu \notin I} \|u_\nu\|^2 \geq \delta$. The boundary of this set is

$$\partial \mathcal{W}_\delta = (\{-\varepsilon\} \times \mathcal{M}(\tau_{-\varepsilon})) \cup (\{\varepsilon\} \times \mathcal{M}(\tau_\varepsilon)) \cup \mathcal{M}_\delta,$$

where the third boundary component $\mathcal{M}_\delta$ is the moduli space of based gauge equivalence classes of solutions of the equations

$$\begin{aligned} \bar{\partial}_A u_\nu &= 0 \quad (\nu = 1, \dots, n), \\ *F_A + \pi \sum_{\nu=1}^n |u_\nu|^2 w_\nu &= \frac{*t\lambda(P)}{\text{Vol}(\Sigma)} + \tau_t, \quad \sum_{\nu \notin I} \|u_\nu\|^2 = \delta. \end{aligned} \tag{13}$$

Here $\|\cdot\|$ denotes the L^2 -norm. If $\delta > 0$ is sufficiently small then $\mathcal{M}_\delta$ does not intersect the boundary of $\mathcal{W}$. We point out that

$$t = \frac{\pi}{\langle \tau_1, e_1 \rangle \text{Vol}(\Sigma)} \sum_{\nu \notin I} \|u_\nu\|_{L^2}^2 \langle w_\nu, e_1 \rangle. \tag{14}$$

To see this take the inner product of the second equation in (13) with e_1 and integrate over Σ . This shows that the parameter t is determined by u and can therefore be removed in the definition of $\mathcal{M}_\delta$. If we orient $\mathcal{M}_\delta$ as the boundary of $\mathcal{W}_\delta$ then the wall crossing number is given by

$$\Phi_{\lambda, g}^{\rho, \tau_0 + \varepsilon \tau_1}(\alpha) - \Phi_{\lambda, g}^{\rho, \tau_0 - \varepsilon \tau_1}(\alpha) = - \int_{\mathcal{M}_\delta / T} \pi^* \alpha.$$

It remains to prove that the integrals over $\mathcal{M}_\delta / T$ and $\mathcal{S}_0 / T$ agree.

We prove this in two steps. The first step shows that the (T -moduli problems associated to) $\mathcal{M}_\delta$ and $\mathcal{S}_0$ are cobordant and the second step is to compare the

orientations. Fix a real parameter $s \in [0, 1]$ and replace the second equation in (13) by

$$*F_A + \pi \sum_{\nu \in I} |u_\nu|^2 w_\nu - \frac{*t\lambda(P)}{\text{Vol}(\Sigma)} - \tau_0 = s \left(t\tau_1 - \pi \sum_{\nu \notin I} |u_\nu|^2 w_\nu \right). \quad (15)$$

For $s = 1$ this is equivalent to the second equation in (13) and for $s = 0$ to the second equation in (12). Thus the moduli spaces $\mathcal{M}_\delta$ and $\mathcal{S}_0$ are equivariantly cobordant and the isotropy subgroups are finite for every s .

Next we compare the orientation of $\mathcal{S}_0/T$ with the boundary orientation of $\mathcal{M}_\delta/T$. Let $[t, A, u] \in \mathcal{W}_\delta$ and define the operator $\mathcal{D} : \mathcal{X} \rightarrow \mathcal{Y}$ by

$$\mathcal{D}(\alpha, \hat{u}) := \begin{pmatrix} *d\alpha + 2\pi \sum_{\nu=1}^n \langle u_\nu, \hat{u}_\nu \rangle w_\nu \\ d^*\alpha + 2\pi \sum_{\nu=1}^n \langle iu_\nu, \hat{u}_\nu \rangle w_\nu \\ \bar{\partial}_A \hat{u}_\nu + \rho_\nu(\alpha)^{0,1} u_\nu \end{pmatrix}.$$

This operator is complex linear. Assume for simplicity that $\mathcal{D}$ is surjective. Then $\mathcal{W}_\delta$ is a manifold near $[t, A, u]$ and the tangent space of $\mathcal{W}_\delta/T$ is

$$T_{[t, A, u]} \mathcal{W}_\delta/T = \{(\hat{t}, \alpha, \hat{u}) \mid \mathcal{D}(\alpha, \hat{u}) = (\hat{t}\tau_1, 0, 0)\}$$

The condition $\mathcal{D}(\alpha, \hat{u}) = (\hat{t}\tau_1, 0, 0)$ implies $\Psi(\alpha, \hat{u}) = \hat{t}$. (Take the inner product of the first two components of $\mathcal{D}(\alpha, \hat{u})$ with e_1 and integrate over Σ .) Thus we may identify

$$T_{[t, A, u]} \mathcal{W}_\delta/T \cong \{(\alpha, \hat{u}) \mid \mathcal{D}(\alpha, \hat{u}) \in (\mathbb{R}\tau_1, 0, 0)\}.$$

Let $v_1, \dots, v_{2m}, w_0$ be a basis of the tangent space with $\text{Re}\Psi(w_0) > 0$ and $v_\nu \in \ker \mathcal{D}$. The basis is called **positively oriented** if the vectors $v_1, \dots, v_{2m}$ form a positively oriented basis of the kernel of $\mathcal{D}$.

Using the determinant bundle over the space of Fredholm operators we can carry the orientation of $\mathcal{W}_\delta$ over to a point $[A, u] \in \mathcal{S}_0$. The notion of an inward pointing vector also carries over. Let $v_1, \dots, v_{2m}$ be a positive basis of $T_{[A, u]} \mathcal{S}_0/T$, consider the inward pointing vector $w_0 := (0, \{u_\nu\}_{\nu \notin I})$, and denote $w_1 := (0, \{-2\pi i \langle w_\nu, e_1 \rangle u_\nu\}_{\nu \notin I})$. Then, by Remark 4.2, the vectors $v_1, \dots, v_{2m}, w_0, w_1$ form a positive basis of $\ker \mathcal{D}_0$. Deform this basis continuously into a basis $\hat{v}_1, \dots, \hat{v}_{2m}, w_0, w_1$ of $\ker \mathcal{D}_0$ such that $\hat{v}_1, \dots, \hat{v}_{2m}$ form a basis of $\ker \mathcal{D}_0 \cap \ker \Psi$. Note that $\text{Im}\Psi(w_1) < 0$. If $\text{Re}\Psi(w_0) < 0$ it follows that the vectors $\hat{v}_1, \dots, \hat{v}_{2m}$ form a positive basis of $\ker \mathcal{D}_0 \cap \ker \Psi$ and otherwise they form a negative basis. Thus, in both cases, the vectors $\hat{v}_1, \dots, \hat{v}_{2m}, w_0$ represent the negative orientation of the tangent space of $\mathcal{W}$. Since w_0 is inward pointing, it follows that the orientation of $\mathcal{S}_0/T$ agrees with the boundary orientation of $\mathcal{M}_\delta/T$. Hence the integral of $\pi^*\alpha$ over $\mathcal{M}_\delta/T$ agrees with the integral over $\mathcal{S}_0/T$.

The assertion about the integrals can be understood verbatim if the moduli spaces are smooth. Otherwise they are understood in terms of the equivariant Euler classes of the associated T -moduli problems. The result then follows from the cobordism axiom for the Euler class in [6]. $\square$

The moduli space $\mathcal{S}_0$ is the sphere bundle in the kernel bundle of a family of Cauchy–Riemann operators over $\mathcal{M}_0$. In Section 4.3 we explain a general equivariant localization formula for such kernel bundles. The relevant index computation uses the Atiyah–Singer index theorem for families and will be carried out in Section 4.4. The next section explains the necessary background about the equivariant Euler class.

4.2 The equivariant Euler class

We begin with some recollections about the equivariant Euler class (see [6] for details). Let X be a compact oriented smooth manifold, $E \rightarrow X$ be an oriented real vector bundle of rank k , and G be a compact Lie group which acts on X and E by orientation preserving diffeomorphisms such that the projection is equivariant and the action is linear on the fibres. We shall think of the action of G on X and E as a right action and denote it by $(x, e) \mapsto (g^*x, g^*e)$ for $e \in E_x$. The corresponding covariant action will be denoted by $g_*x := (g^{-1})^*x$ and the infinitesimal (contravariant) action of $\xi \in \mathfrak{g} := \text{Lie}(G)$ by $\xi^*x \in T_xX$. An **equivariant Thom form** is a d_G -closed equivariant differential form $\tau_G(E) \in \Omega_G^k(E)$ with compact support and fibre integral one. The **equivariant Euler class** $e_G(E) \in H_G^k(X)$ is the cohomology class of the pullback of an equivariant Thom form under the zero section. We will sometimes use the same notation for the Euler class and a form representing it. The Thom class and the Euler class are multiplicative under direct sum.

Now suppose that E is a rank n complex vector bundle and the action of G is complex linear on the fibres. Then an explicit representative of the equivariant Euler class can be constructed as follows. Fix a G -invariant Hermitian metric on E and let $P \rightarrow X$ denote the unitary frame bundle of E . This bundle carries a right action of $U(n)$ and a left action of G , and these two actions commute. Let $X_\xi \in \text{Vect}(P)$ denote the infinitesimal (covariant) action of $\xi \in \mathfrak{g}$. More precisely a point $p \in P_x$ of the fibre over $x \in X$ is a unitary vector space isomorphism $p: \mathbb{C}^n \rightarrow E_x$ and the left action of $g \in G$ is given by $g_*p: \mathbb{C}^n \rightarrow E_{g_*x}$. The vector field $X_\xi \in \text{Vect}(P)$ is defined by

$$X_\xi(p) := \left. \frac{d}{dt} \right|_{t=0} \exp(t\xi)_*p \in T_pP.$$

Lemma 4.3. *Let $A \in \mathcal{A}(P) \subset \Omega^1(P, \mathfrak{u}(n))$ be a G -invariant $U(n)$ -connection form on P . Then the G -equivariant Euler class of a complex vector bundle E is represented by the d_G -closed form*

$$e_G(E, \xi) = \det \left(\frac{i}{2\pi} F_A + \frac{i}{2\pi} A(X_\xi) \right), \quad (16)$$

where $F_A \in \Omega^2(P, \mathfrak{u}(n))$ denotes the curvature of A .

Proof. The right hand side in (16) is invariant and horizontal for the $U(n)$ -action and thus descends to a G -equivariant form on X . It is easy to check that this form is d_G -closed and hence represents an equivariant cohomology class.

To prove (16) we assume first that $E = X \times \mathbb{C}^n$ is a trivial bundle and $\rho : G \rightarrow U(n)$ is a unitary representation of G . The homomorphism ρ defines the covariant action of G on E and so

$$g^*(x, z) = (g^*x, \rho(g)^{-1}z)$$

for $x \in X$, $z \in \mathbb{C}^n$, and $g \in G$. The frame bundle of E is the product bundle $P := X \times U(n)$ and the formula

$$A_{x,u}(v, u\eta) := \eta$$

for $v \in T_x X$, $u \in U(n)$ and $\eta \in \mathfrak{u}(n)$ defines a $U(n)$ -connection form $A \in \Omega^1(P, \mathfrak{u}(n))$. This connection is G -invariant and flat. For $\xi \in \mathfrak{g}$ the vector field $X_\xi \in \text{Vect}(P)$ is given by

$$X_\xi(x, u) = (0, \dot{\rho}(\xi)u)$$

and so

$$A(X_\xi(x, u)) = u^{-1}\dot{\rho}(\xi)u, \quad \det \left(\frac{i}{2\pi} A(X_\xi) \right) = \det \left(\frac{i}{2\pi} \dot{\rho}(\xi) \right).$$

Now a Thom form on E can be constructed as follows. For $k = 0, \dots, n$ let $\sigma_k : \mathfrak{u}(n) \rightarrow \Omega^{2n-2k}(\mathbb{C}^n)$ be a polynomial map of degree k . It is shown in [6, Lemma 5.5] that these polynomials can be chosen such that $\sigma_0 \in \Omega^{2n}(\mathbb{C}^n)$ is the standard volume form,

$$\sigma_n(\eta) = \det(i\eta),$$

and

$$\iota(v_\eta)\sigma_k(\eta) = \lambda \wedge \sigma_{k+1}(\eta)$$

for each k , where $\lambda \in \Omega^1(\mathbb{C}^n)$ is the differential of the function $z \mapsto |z|^2/2$ and the vector field $v_\eta \in \text{Vect}(\mathbb{C}^n)$ is defined by $v_\eta(z) := \eta z$ for $\eta \in \mathfrak{u}(n)$. Now choose functions $f_k : [0, \infty) \rightarrow [0, \infty)$ with compact support such that $f_0(s) = 0$ for $s \leq \delta$ and $s \geq 1$ and

$$f'_k(s) + f_{k-1}(s) = 0, \quad f_k(1) = 0,$$

and

$$\int_0^\infty s^k f_0(s) ds = 0, \quad \int_0^\infty s^{n-1} f_0(s) ds = \frac{1}{2^{n-1} \text{Vol}(S^{2n-1})}$$

for $0 \leq k \leq n-2$. Then

$$f_k(s) = \frac{1}{(k-1)!} \int_s^1 (t-s)^{k-1} f_0(t) dt$$

and hence $f_k(s) = 0$ for $s < \delta$ and $k < n$ and

$$f_n(0) = \frac{1}{2^{n-1}(n-1)! \text{Vol}(S^{2n-1})} = \frac{1}{(2\pi)^n}.$$

Now a Thom form on $E = X \times \mathbb{C}^n$ is given by

$$\tau(\xi) = \sum_{k=1}^n f_k(|z|^2/2) \sigma_k(\dot{\rho}(\xi)).$$

Its pullback under the zero section is given by

$$e_G(E) = f_n(0) \sigma_n(\dot{\rho}(\xi)) = \det \left(\frac{i}{2\pi} \dot{\rho}(\xi) \right).$$

This proves the lemma in the case $E = X \times \mathbb{C}^n$. For general G -equivariant bundles $E \rightarrow X$ the result follows from the (*Naturality*) axiom for the Euler class and the fact that the pullback of E under the projection $P \rightarrow E$ is isomorphic to the trivial bundle $P \times \mathbb{C}^n$. $\square$

Remark 4.4. The formula of Lemma 4.3 can also be expressed as follows. Let ∇ be a G -equivariant Hermitian connection on E and, for $\xi \in \mathfrak{g}$, denote by $\xi^\nabla \in \Omega^0(X, \text{End}(E))$ the covariant infinitesimal action defined by

$$\xi^\nabla e := \nabla_t \exp(t\xi)_* e|_{t=0}$$

Then the Euler class is given by

$$e_G(E, \xi) = \det \left(\frac{i}{2\pi} F^\nabla + \frac{i}{2\pi} \xi^\nabla \right).$$

Example 4.5. Let $E \rightarrow X$ be a rank n complex vector bundle. Suppose G acts trivially on X and that the covariant action on the fibres is given by a homomorphism $\rho : G \rightarrow S^1$, given by

$$\rho(\exp(\xi)) = e^{-2\pi i \langle w, \xi \rangle},$$

where $w \in \mathfrak{g}^*$. Then, for every G -invariant Hermitian connection ∇ on E , the endomorphism $\xi^\nabla \in \Omega^0(X, \text{End}(E))$ is given by multiplication with the imaginary number $\dot{\rho}(\xi)$. Hence

$$e_G(E, \xi) = \det \left(\frac{i}{2\pi} F^\nabla + \frac{i}{2\pi} \dot{\rho}(\xi) \right) = \sum_{j=0}^n \langle w, \xi \rangle^{n-j} c_j(E).$$

We wish to invert the equivariant Euler class. This requires an extension of the equivariant cohomology ring of X .

Standing assumption. *In the following X is a smooth manifold, G is a compact Lie group acting on X , and $T_1 \subset G$ is an oriented circle which is contained in the center of G and acts trivially on X .*

Denote the quotient group by $G_0 := G/T_1$. Denote by e_1 the positive integral generator of the Lie algebra $\mathfrak{t}_1 := \text{Lie}(T_1)$. Let n be an integer. A T_1 -**rational**

G-equivariant differential form of degree n on X is a Laurent series in z^{-1} of the form

$$\alpha(\xi, z) = \sum_{j \leq n/2} \alpha_j(\xi) z^j,$$

with coefficients $\alpha_j \in \Omega_{\mathbf{G}}^{n-2j}(X)$, that satisfies the following conditions.

(i) For every $\xi \in \mathfrak{g}$ and every $x \in X$ the Laurent series $\sum_{j \leq n/2} \alpha_j(\xi)_x z^j$ is a rational function on $\mathbb{C}$ with values in the complex vector space $\Lambda^* T_x^* X \otimes \mathbb{C}$.

(ii) For every $t \in \mathbb{R}$ we have $\alpha(\xi, t + z) = \alpha(\xi + te_1, z)$. Equivalently,

$$\alpha_k(\xi + te_1) = \sum_{k \leq j \leq n/2} \binom{j}{k} \alpha_j(\xi) t^{j-k}, \quad k \geq 0, \quad (17)$$

$$\alpha_k(\xi + te_1) = \sum_{k \leq j < 0} \binom{-k-1}{-j-1} \alpha_j(\xi) (-t)^{j-k}, \quad k < 0. \quad (18)$$

Denote by $\Omega_{\mathbf{G}, T_1}^n(X)$ the space of T_1 -rational G-equivariant differential forms on X . This is a chain complex with respect to the usual equivariant differential $d_{\mathbf{G}} \alpha(\xi) := d\alpha(\xi) + \iota(X_{\xi})\alpha(\xi)$. The cohomology of this chain complex will be denoted by $H_{\mathbf{G}, T_1}^*(X)$.

Let $\alpha = \sum_j \alpha_j z^j \in \Omega_{\mathbf{G}, T_1}^n(X)$. Then $\alpha_{-1}(\xi + te_1) = \alpha_{-1}(\xi)$. In other words, the coefficient of z^{-1} descends to a $\mathbf{G}_0$ -equivariant cohomology class on X . Minus this coefficient is called **the residue at infinity of α** and will be denoted by

$$\text{Res}_{\infty}(\alpha) := -\alpha_{-1}(\xi) = -\frac{1}{2\pi i} \oint \alpha(\xi, z) dz \in \Omega_{\mathbf{G}_0}^{n+2}(X).$$

The residue at infinity descends to a homomorphism

$$\text{Res}_{\infty} : H_{\mathbf{G}, T_1}^n(X) \rightarrow H_{\mathbf{G}_0}^{n+2}(X).$$

Remark 4.6. There is an obvious inclusion $\Omega_{\mathbf{G}}^*(X) \rightarrow \Omega_{\mathbf{G}, T_1}^*(X)$ whose image is the subspace of polynomials $\alpha = \sum_{0 \leq j \leq n/2} \alpha_j z^j \in \Omega_{\mathbf{G}, T_1}^n(X)$. Condition (17) shows that any such form is uniquely determined by $\alpha_0 \in \Omega_{\mathbf{G}}^n(X)$ and vice versa. The inclusion $\Omega_{\mathbf{G}}^*(X) \hookrightarrow \Omega_{\mathbf{G}, T_1}^*(X)$ induces an inclusion in cohomology

$$H_{\mathbf{G}}^*(X) \hookrightarrow H_{\mathbf{G}, T_1}^*(X)$$

whose left inverse is induced by the projection $\alpha = \sum_j \alpha_j z^j \mapsto \alpha_0$.

Let $E \rightarrow X$ be a G-equivariant complex vector bundle of rank n_E . The subgroup T_1 acts on E with **weight**

$$\mathbf{w}_E := \det \left(\frac{i}{2\pi} \dot{\rho}_x(e_1) \right) \in \mathbb{Z}.$$

Here the homomorphism $\rho_x : T_1 \rightarrow \text{Aut}(E_x)$ denotes the action on the fiber over x and $\dot{\rho}_x : \mathfrak{t}_1 \rightarrow \text{End}(E_x)$ denotes the corresponding Lie algebra homomorphism. The weight w_E is independent of x . Think of the equivariant Euler class as a polynomial map $\mathfrak{g} \rightarrow \Omega^*(X)$. By Lemma 4.3, the G-equivariant Chern classes $c_j(E) \in H_G^{2j}(X)$ are the coefficients of z^{n_E-j} in the polynomial

$$e_G(E, \xi + ze_1) =: \sum_{j=0}^{n_E} c_j(E, \xi) z^{n_E-j}.$$

In particular, $c_0(E, \xi) = w_E$. If $w_E \neq 0$ then the equivariant Euler class $e_G \in H_T^{2n_E}(X)$ has a well defined inverse $1/e_G$ in the T_1 -rational G-equivariant cohomology group $H_{G, T_1}^{-2n_E}(X)$. To see this, expand the rational function $z \mapsto 1/e_G(\xi + ze_1)$ into a Laurent series in z^{-1} which converges near infinity:

$$\begin{aligned} \frac{1}{e_G(E, \xi + ze_1)} &= \frac{1}{w_E z^{n_E}} \sum_{k=0}^{\infty} \left(- \sum_{j=1}^{n_E} \frac{c_j(E, \xi)}{w_E} z^{-j} \right)^k \\ &=: \sum_{i=0}^{\infty} s_i(E, \xi) z^{-n_E-i}. \end{aligned}$$

The coefficients $s_i(E) \in H_G^{2i}(X)$ of this Laurent series are called the **equivariant Segre classes** of E . They are uniquely determined by the equation

$$\sum_{i+j=k} s_i(E, \xi) c_j(E, \xi) = \begin{cases} 1, & \text{if } k = 0, \\ 0, & \text{if } k > 0. \end{cases} \quad (19)$$

In particular, the degree zero Segre class is $s_0(E, \xi) = 1/w_E$. If $F \rightarrow X$ is another G-equivariant complex vector bundle of rank n_F with weight w_F , then the quotient

$$e_G(F \ominus E) := \frac{e_G(F)}{e_G(E)} \in H_{G, T_1}^{2n_F-2n_E}(X)$$

depends only on the equivariant K -theory class $F \ominus E \in K_G(X)$. It is only defined for equivariant K -theory classes $F \ominus E$ whose denominator E has nonzero weight.

4.3 Localization

Let X be an orientable smooth manifold, $G = T$ be a torus acting on X , and $T_1 \subset T$ be an oriented circle that acts trivially on X . We assume that the quotient group $T_0 := T/T_1$ acts on X with finite isotropy. Denote the Lie algebras by $\mathfrak{t} := \text{Lie}(T)$, $\mathfrak{t}_1 := \text{Lie}(T_1)$, and $\mathfrak{t}_0 := \mathfrak{t}/\mathfrak{t}_1 := \text{Lie}(T_0)$, let $\Lambda \subset \mathfrak{t}$ be the integer lattice, and denote by $e_1 \in \mathfrak{t}_1 \cap \Lambda$ the positive generator of the sublattice. Throughout we denote $m := \dim X - \dim T_0$.

Let $\mathcal{E} \rightarrow X$ and $\mathcal{F} \rightarrow X$ be complex Hilbert space bundles on which T acts complex linearly such that the projections are equivariant. Assume that T_1 acts with finite isotropy outside the zero sections of $\mathcal{E}$ and $\mathcal{F}$. Let

$$\mathcal{D}_x : \mathcal{E}_x \rightarrow \mathcal{F}_x$$

be a smooth family of G -equivariant complex linear Fredholm operators of complex numerical index

$$\text{index}(\mathcal{D}) := \dim^{\mathbb{C}} \ker \mathcal{D}_x - \dim^{\mathbb{C}} \text{coker } \mathcal{D}_x.$$

Denote by

$$\mathcal{IND}(\mathcal{D}) := \bigcup_{x \in X} \{x\} \times \ker \mathcal{D}_x \ominus \text{coker } \mathcal{D}_x \in K_G(X)$$

the topological index of $\mathcal{D}$, understood as a G -equivariant K -theory class. Consider the following G -moduli problem. The Hilbert manifold $\mathbb{B}$ is given by

$$\mathbb{B} := \left\{ (x, e) \mid x \in X, e \in \mathcal{E}_x, \|e\|^2 = 1 \right\},$$

the Hilbert space bundle $\mathbb{H} \rightarrow \mathbb{B}$ has fibre

$$\mathbb{H}_{x,e} := \mathcal{F}_x$$

over $(x, e) \in \mathbb{B}$, and the section $\mathbb{S} : \mathbb{B} \rightarrow \mathbb{H}$ is given by

$$\mathbb{S}(x, e) := \mathcal{D}_x e.$$

The zero set of this section is the *kernel manifold*

$$\mathbb{M} := \{(x, e) \in \mathbb{B} \mid \mathcal{D}_x e = 0\}.$$

Denote by $\pi : \mathbb{B} \rightarrow X$ the obvious projection. The equivariant K -theory class $\mathcal{IND}(\mathcal{D}) \in K_G(X)$ has a nonzero weight (for the T_1 -action) and hence carries an equivariant Euler class

$$e_G(\mathcal{IND}(\mathcal{D})) \in H_{T, T_1}^*(X).$$

in the T_1 -rational T -equivariant cohomology of X . The following theorem generalizes the localization formula for circle actions in [6]. The assertion requires a choice of orientations.

Remark 4.7. Orientations of X/T_0 and T_1 determine an orientation of the T -moduli problem $(\mathbb{B}, \mathbb{H}, \mathbb{S})$ as follows. By choosing local trivializations we may assume that $\mathcal{E}$ and $\mathcal{F}$ are (complex) Hilbert spaces equipped with a T -action and so $\mathcal{D}$ is a T -equivariant smooth map $X \rightarrow \mathcal{L}(\mathcal{E}, \mathcal{F}) : x \mapsto \mathcal{D}_x$, which assigns a (complex linear) Fredholm operator $\mathcal{D}_x$ to every $x \in X$. In this case the vertical differential of $\mathbb{S}$ at a point $(x, e) \in \mathbb{M}$ is an operator

$$D\mathbb{S}(x, e) : \{(\hat{x}, \hat{e}) \in T_x X \times \mathcal{E} \mid \langle \hat{e}, e \rangle = 0\} \rightarrow \mathcal{F}.$$

It is given by

$$D\mathbb{S}(x, e)(\hat{x}, \hat{e}) = \mathcal{D}_x \hat{e} + \dot{\mathcal{D}}(\hat{x})e,$$

where $\dot{\mathcal{D}}(\hat{x})e$ is defined as the derivative of the path $\mathbb{R} \rightarrow \mathcal{F} : t \mapsto \mathcal{D}_{\exp_x(t\hat{x})}e$ at $t = 0$. Now suppose that $\mathcal{D}_x$ is surjective. Then a **positive basis** of the kernel of $D\mathbb{S}(x, e)$ is defined as follows. Pick a positive basis $\hat{x}_1, \dots, \hat{x}_m$ of $T_x X / \mathfrak{t}_0 x$ and choose $\hat{e}_1, \dots, \hat{e}_m \in \mathcal{E}$ such that $\mathcal{D}_x \hat{e}_i + \dot{\mathcal{D}}(\hat{x}_i)e = 0$ and $\langle \hat{e}_i, e \rangle = 0$ for $i = 1, \dots, m$. Next choose a positive basis $\hat{e}_{m+1}, \dots, \hat{e}_{m+2n}$ of the complex vector space $\ker \mathcal{D}_x$ such that $\hat{e}_{m+2n-1} = e$ and $\hat{e}_{m+2n}$ is a positive tangent vector of the T_1 -orbit of e . Then the vectors $(\hat{x}_i, \hat{e}_i)$ for $i = 1, \dots, m$ and the vectors $(0, \hat{e}_j)$ for $j = m+1, \dots, m+2n-2$ are declared to be a positive basis of $\ker D\mathbb{S}(x, e) / \mathfrak{t} \cdot (x, e)$. This definition of the orientation is independent of the choices. If $\mathcal{D}_x$ is not surjective, one can apply the same construction to the kernel of a suitably augmented operator. We emphasize that the orientation described here agrees with the convention of Remark 4.2.

Theorem 4.8. *Let $m := \dim X - \dim T_0$ and $n := \text{index}(\mathcal{D})$. Fix any orientation of X/T_0 , let T_1 be oriented by e_1 , and orient $\mathbb{M}/T$ as in Remark 4.7. Then*

$$\int_{\mathbb{M}/T} \pi^* \alpha = \int_{X/T_0} \text{Res}_\infty \left(\frac{\alpha}{e_T(\mathcal{I}\mathcal{N}\mathcal{D}(\mathcal{D}))} \right) \quad (20)$$

for every $\alpha \in H_T^{m+2n-2}(X)$.

The integral on the left is understood as the Euler class of the T -moduli problem $(\mathbb{B}, \mathbb{H}, \mathbb{S})$ evaluated on $\pi^* \alpha$ (see [6]). The integrand on the right is the residue at infinity of the T_1 -rational T -equivariant cohomology class $\alpha / e_T(\mathcal{I}\mathcal{N}\mathcal{D}(\mathcal{D})) \in H_{T, T_1}^{m-2}(X)$. It is a T_0 equivariant cohomology class in $H_{T_0}^m(X)$ and can be integrated over X/T_0 because T_0 acts on X with finite isotropy.

Remark 4.9. Theorem 4.8 continues to hold if we replace X by a T_0 -moduli problem $(\mathcal{B}_0, \mathcal{E}_0, \mathcal{S}_0)$ as in [6] and $\mathcal{E}$ and $\mathcal{F}$ by Hilbert space bundles over $\mathcal{B}_0$. Then $\mathbb{B}$ is the unit sphere bundle in $\mathcal{E}$, $\mathbb{H}_{b,e} = \mathcal{E}_{0b} \oplus \mathcal{F}_b$, $\mathbb{S}(b, e) = (\mathcal{S}_0(b), \mathcal{D}_b e)$, and the right hand side of (20) is understood in terms of the Euler class of $(\mathcal{B}_0, \mathcal{E}_0, \mathcal{S}_0)$. To prove this, choose a finite dimensional reduction of $(\mathcal{B}_0, \mathcal{E}_0, \mathcal{S}_0)$ and note that (20) continues to hold for noncompact manifolds X and compactly supported T -equivariant differential forms α .

Proof of Theorem 4.8. The proof has three steps.

Step 1. *We may assume without loss of generality that $\mathcal{E}$ is finite dimensional and admits an equivariant trivialization and that $\mathcal{F} = 0$.*

The reduction to the finite dimensional case is proved as in [6, Theorem 11.1]. Hence assume $E = \mathcal{E}$ and $F = \mathcal{F}$ are finite dimensional. By Proposition A.1, there exists a T -equivariant complex vector bundle $E' \rightarrow X$ such that $E \oplus E'$ is equivariantly isomorphic to $X \times V$ for some complex T -representation V . Since T is a torus there exists a homomorphism $T \rightarrow S^1$ whose restriction to T_1 has nonzero degree. Multiplying the action of T on E' with a suitable power of this

homomorphism we may assume that the action of T_1 on E' has nonzero weight. Now let $\mathbb{B}' \subset E \oplus E'$ be the unit sphere bundle, $\mathbb{H}' \rightarrow \mathbb{B}'$ be the pullback of $F \oplus E'$ under the projection $\pi' : \mathbb{B}' \rightarrow X$, and $\mathbb{S}' : \mathbb{B}' \rightarrow \mathbb{H}'$ be given by

$$\mathbb{S}'(x, e, e') := (x, D_x e, e').$$

Then the inclusion $\mathbb{B} \rightarrow \mathbb{B}' : (x, e) \mapsto (x, e, 0)$ defines a morphism of T -moduli problems. Hence, assuming the assertion for E replaced by $X \times V$ and F replaced by the zero bundle, we obtain

$$\begin{aligned} \int_{\mathbb{M}/T} \pi^* \alpha &= \chi^{\mathbb{B}, \mathbb{H}, \mathbb{S}}(\pi^* \alpha) \\ &= \chi^{\mathbb{B}', \mathbb{H}', \mathbb{S}'}(\pi'^* \alpha) \\ &= \int_{\mathbb{B}'/T} \pi'^* \alpha \wedge e_T(\mathbb{H}') \\ &= \int_{\mathbb{B}'/T} \pi'^* \alpha \wedge \pi'^* e_T(F \oplus E') \\ &= \int_{X/T_0} \text{Res}_\infty \left(\frac{\alpha \wedge e_T(F \oplus E')}{e_T(E \oplus E')} \right) \\ &= \int_{X/T_0} \text{Res}_\infty \left(\frac{\alpha}{e_T(\mathcal{IN}\mathcal{D}(D))} \right). \end{aligned}$$

Here the second equation uses the (*Naturality*) axiom for the Euler class, the third equation uses the (*Thom class*) axiom (see [6]), the fourth equation uses the fact that $\mathbb{H}'$ is the pullback of $F \oplus E'$, and the fifth equation uses the hypothesis that the result holds when E is a trivial bundle and $F = 0$.

Step 2. Suppose $E = L = X \times \mathbb{C}$ is a trivial line bundle and denote by $B \subset L$ the unit circle bundle. Then for every $\alpha = \sum_{j \leq m/2} \alpha_j z^j \in \Omega_{T, T_1}^m(X)$,

$$\int_{B/T} \pi^* \alpha_0 = \int_{X/T_0} \text{Res}_\infty \left(\frac{\alpha}{e_T(L)} \right). \quad (21)$$

Let $\rho : T \rightarrow S^1$ denote the covariant action of T on the fibres of L and suppose that T_1 acts on the fibers with weight ℓ . Then $\dot{\rho}(e_1) = -2\pi i \ell$ and, by Lemma 4.3,

$$e_T(L, \xi + z e_1) = \ell z + \frac{i \dot{\rho}(\xi)}{2\pi}.$$

Each form $\alpha_j \in \Omega_T^{m-2j}(X)$ is equivariantly closed and hence represents a T -equivariant cohomology class on X . Now

$$\begin{aligned} \frac{\alpha(\xi, z)}{e_T(\xi + z e_1)} &= \left(\sum_{j \leq m/2} \frac{\alpha_j(\xi)}{\ell} z^{j-1} \right) \frac{1}{1 + \frac{i \dot{\rho}(\xi)}{2\pi \ell z}} \\ &= \left(\sum_{j \leq m/2} \frac{\alpha_j(\xi)}{\ell} z^{j-1} \right) \sum_{k \geq 0} \left(-\frac{i \dot{\rho}(\xi)}{2\pi \ell z} \right)^k. \end{aligned}$$

The residue at infinity is minus the coefficient of z^{-1} in this power series. Thus

$$\text{Res}_\infty \left(\frac{\alpha}{e_T(L)} \right) = -\frac{1}{\ell} \sum_{j \geq 0} \alpha_j(\xi) \left(-\frac{i\dot{\rho}(\xi)}{2\pi\ell} \right)^j \quad (22)$$

By (17), the right hand side is invariant under the shift $\xi \mapsto \xi + te_1$ and hence descends to a T_0 -equivariant differential form on X .

To compute the integral of $\pi^*\alpha_0$ over B/T , we denote the elements of $B = X \times S^1$ by (x, u) , where $x \in X$ and $u \in S^1$. Then $u^{-1}du$ is the standard $U(1)$ -connection form on B . Define $A_1 \in \Omega^1(B, \mathfrak{t}_1)$ by

$$A_1 := -\frac{i u^{-1} du}{2\pi\ell} e_1.$$

This form is T -invariant and satisfies $A_1(0, -\dot{\rho}(e_1)z) = (i\dot{\rho}(e_1)/2\pi\ell)e_1 = e_1$. Hence it is a T -invariant T_1 -connection as in [6], where we regard the action by ρ^{-1} as the contravariant action on B . The infinitesimal covariant action of $\xi \in \mathfrak{t}$ on B is given by $X_\xi(x, u) = (0, \dot{\rho}(\xi)u)$. Hence the T -equivariant curvature of A_1 is the 2-form $F_{A_1, T} \in \Omega_T^2(B, \mathfrak{t})$ given by

$$F_{A_1, T}(\xi) := F_{A_1} + \xi + A_1(X_\xi) = \xi - \frac{i\dot{\rho}(\xi)}{2\pi\ell} e_1.$$

(see [6, Section 3]). Replacing ξ by $F_{A_1, T}(\xi)$ in the equivariant differential form $\pi^*\alpha_0 \in \Omega_T^m(B)$ we obtain the T_1 -basic T -equivariant differential form

$$(\pi^*\alpha_0)_{A_1} = \pi^*\alpha_0 \left(\xi - \frac{i\dot{\rho}(\xi)}{2\pi\ell} e_1 \right) = \sum_{j \geq 0} \pi^*\alpha_j(\xi) \left(-\frac{i\dot{\rho}(\xi)}{2\pi\ell} \right)^j$$

on $B = X \times S^1$. The projection $\pi : B \rightarrow X$ induces a T_0 -equivariant diffeomorphism from B/T_1 to X , however, each point in B has an isotropy subgroup of order ℓ under the action of T_1 . Moreover, the diffeomorphism is orientation preserving if and only if ℓ is negative. (If $\xi_1, \dots, \xi_m$ is a positive basis of $T_x X/T_0$ and $u \in S^1$ then, according to Remark 4.7, the basis $(\xi_1, 0), \dots, (\xi_m, 0)$ of $T_{(x, u)} B/T$ is positive if and only if the vectors $(\xi_1, 0), \dots, (\xi_m, 0), (0, u), (0, \dot{\rho}(e_1)u)$ form a positive basis of $T_x X/T_0 \times \mathbb{C}$. Since $\dot{\rho}(e_1) = -2\pi i\ell$, this is the case if and only if ℓ is negative.) Hence

$$\int_{B/T} \pi^*\alpha_0 = \int_{B/T} (\pi^*\alpha_0)_{A_1} = -\frac{1}{\ell} \sum_{j \geq 0} \int_{X/T_0} \alpha_j(\xi_0) \left(-\frac{i\dot{\rho}(\xi_0)}{2\pi\ell} \right)^j,$$

and so the assertion of Step 2 follows from (22).

Step 3. *We prove the theorem.*

By Step 1, we may assume without loss of generality that $F = 0$ and $E = X \times V$ for some unitary T -representation V . Since T is a torus, we may assume that

$V = \mathbb{C}^n$ and that T acts diagonally by homomorphisms $\rho_\nu : T \rightarrow S^1$ for $\nu = 1, \dots, n$. Denote by $L_\nu := X \times \mathbb{C}$ the T -equivariant bundle where T acts by ρ_ν on the fibre. Consider the T -moduli problem (B, H, S) given by

$$B := X \times S^{2n-1}, \quad H := \pi^* L_1 \oplus \dots \oplus \pi^* L_{n-1},$$

where $\pi : B \rightarrow X$ denotes the projection, and

$$S(x, z_1, \dots, z_n) := (z_1, \dots, z_{n-1}).$$

Then the T -equivariant Euler class of H is the pullback under π of the Euler class of $L_1 \oplus \dots \oplus L_{n-1}$, i.e.

$$e_T(H) = \pi^* e_T(L_1) \wedge \dots \wedge \pi^* e_T(L_{n-1}).$$

Let $\alpha \in \Omega_T^{m+2n-2}(X)$ and define $\beta \in \Omega_{T, T_1}^m(X)$ by

$$\beta := \sum_{j \leq m/2} \beta_j z^j := \frac{\alpha}{e_T(L_1) \dots e_T(L_{n-1})}, \quad \beta_j \in \Omega_T^{m-2j}(X).$$

Since T acts on B with finite isotropy, we can represent the equivariant cohomology class $e_T(H) \in H_T^{2n-2}(B)$ by a T -invariant and horizontal differential form $\tau_H \in \Omega^{2n-2}(B)$ (see [6, Theorem 3.8]). With such a representative the identity $\pi^* \alpha = \pi^* \beta \wedge e_T(H)$ in $\Omega_{T, T_1}^*(X)$ takes the form $\pi^* \alpha = \pi^* \beta_0 \wedge \tau_H$. Now S is transverse to the zero section and $S^{-1}(0)$ is the unit sphere bundle $B_n := B \cap L_n$ in L_n . Hence it follows from the (*Transversality*) axiom for the Euler class in [6] that

$$\int_{B/T} \pi^* \alpha = \int_{B/T} \pi^* \beta_0 \wedge \tau_H = \int_{B_n/T} \pi^* \beta_0 = \int_{X/T_0} \text{Res}_\infty \left(\frac{\beta}{e_T(L_n)} \right).$$

The last equation follows from Step 2. Since $\beta/e_T(L_n) = \alpha/e_T(E)$, this proves the theorem. $\square$

4.4 The index formula

We return to the setting of Section 3. Recall that $P \rightarrow \Sigma$ is a principal T -bundle and $L_\nu = P \times_{\rho_\nu} \mathbb{C} \rightarrow \Sigma$ for $\nu = 1, \dots, n$. Given an index set $I \subset \{1, \dots, n\}$ as in Section 3 we consider the principal $\mathcal{G}_0$ -bundle

$$\mathcal{P}_I := \mathcal{A}(P) \times \bigoplus_{\nu \in I} \Omega^0(\Sigma, L_\nu) \rightarrow \mathcal{B}_I := \mathcal{P}_I / \mathcal{G}_0$$

where the based gauge group $\mathcal{G}_0$ acts by $g^*(A, u_\nu) = (A + g^{-1}dg, \rho_\nu(g)^{-1}u_\nu)$. It also acts on L_{ν_0} by $g^*[p, \zeta] := [pg(z)^{-1}, \zeta] = [p, \rho_{\nu_0}(g(z))^{-1}\zeta]$, where $z := \pi(p) \in \Sigma$. For $\nu_0 \notin I$ we consider the universal line bundle

$$\mathbb{L}^{\nu_0} := \frac{\mathcal{P}_I \times L_{\nu_0}}{\mathcal{G}_0} \rightarrow \mathcal{B}_I \times \Sigma.$$

The torus T acts on $\mathbb{L}^{\nu_0}$ by $h^*[A, u_\nu, p, \zeta] := [A, \rho_\nu(h)^{-1}u_\nu, p, \rho_{\nu_0}(h)^{-1}\zeta]$.

For $\mathbf{x} = [A, u] \in \mathcal{B}_I$ let us denote by $\mathbb{L}_\mathbf{x}^\nu$ the restriction of $\mathbb{L}^\nu$ to $\{\mathbf{x}\} \times \Sigma$. This restriction is equipped with a natural connection (induced by A) and hence with a Cauchy-Riemann operator

$$\bar{\partial}_\mathbf{x}^\nu : \Omega^0(\Sigma, \mathbb{L}_\mathbf{x}^\nu) \rightarrow \Omega^{0,1}(\Sigma, \mathbb{L}_\mathbf{x}^\nu).$$

Next consider the universal vector bundle

$$\mathbb{E} := \bigoplus_{\nu \notin I} \mathbb{L}^\nu$$

and its restrictions $\mathbb{E}_\mathbf{x}$ to $\{\mathbf{x}\} \times \Sigma$. The Cauchy-Riemann operators

$$\bar{\partial}_\mathbf{x} : \Omega^0(\Sigma, \mathbb{E}_\mathbf{x}) \rightarrow \Omega^{0,1}(\Sigma, \mathbb{E}_\mathbf{x})$$

form a family of Fredholm operators over $\mathcal{B}_I$ between appropriate Hilbert space completions $\mathcal{E}_\mathbf{x}$ of $\Omega^0(\Sigma, \mathbb{E}_\mathbf{x})$ and $\mathcal{F}_\mathbf{x}$ of $\Omega^{0,1}(\Sigma, \mathbb{E}_\mathbf{x})$. These operators are complex linear and equivariant with respect to the action of T .

As in Section 3 we denote by T_1 the identity component of the isotropy subgroup of the subspace $\mathbb{C}^I := \{x \in \mathbb{C}^n \mid x_\nu = 0 \text{ for } \nu \notin I\}$ and assume that T_1 is a circle. This circle acts trivially on the base $\mathcal{B}_I$ and with finite isotropy outside of the zero sections of $\mathcal{E}$ and $\mathcal{F}$. The quotient group $T_0 := T/T_1$ acts with finite isotropy on the moduli space $\mathcal{M}_0 \subset \mathcal{B}_I$ of solutions of equation (8). Hence we are in the situation of Theorem 4.8. The relevant dimensions are

$$\begin{aligned} \dim \mathcal{M} - \dim T &= (n - \dim T)(2 - 2g) + 2 \sum_{\nu=1}^n d_\nu =: 2m, \\ \dim \mathcal{M}_0 - \dim T_0 &= (|I| - \dim T_0)(2 - 2g) + 2 \sum_{\nu \in I} d_\nu + 2g, \\ \text{index}(\bar{\partial}) &= \sum_{\nu \notin I} (d_\nu + 1 - g), \end{aligned}$$

where d_ν denotes the first Chern number of the bundle $L_\nu \rightarrow \Sigma$. Note that the kernel manifold of $\bar{\partial}$ is precisely the space $\mathcal{S}_0$ of solutions of equations (12). Hence, by Theorem 4.8 and Remark 4.9, we have

$$\begin{aligned} \int_{\mathcal{S}_0/T} \pi^* \alpha &= \int_{\mathcal{M}_0/T_0} \text{Res}_\infty \left(\frac{\alpha}{e_T(\mathcal{IND}(\bar{\partial}))} \right) \\ &= \int_{\mathcal{M}_0/T_0} \text{Res}_\infty \left(\frac{\alpha}{\prod_{\nu \notin I} e_T(\mathcal{IND}(\bar{\partial}^\nu))} \right). \end{aligned} \quad (23)$$

Lemma 4.10. *Denote by $\pi_0^* : H_{T, T_1}^*(\mathcal{A}/\mathcal{G}_0) \rightarrow H_{T, T_1}^*(\mathcal{B}_I)$ the homomorphism induced by the projection $\pi_0 : \mathcal{B}_I \rightarrow \mathcal{A}/\mathcal{G}_0$. Then, for every $\nu \notin I$,*

$$e_T(\mathcal{IND}(\bar{\partial}^\nu)) = \pi_0^* \langle \mathbf{w}_\nu, \xi \rangle^{d_\nu + 1 - g} \exp \left(- \frac{\Omega_\nu}{\langle \mathbf{w}_\nu, \xi \rangle} \right) \in H_{T, T_1}^*(\mathcal{B}_I).$$

Proof of Theorem 3.1. By Proposition 4.1 and (23) we have

$$\begin{aligned}
\Phi_{\lambda,g}^{\rho,\tau_0+\varepsilon\tau_1}(\alpha) - \Phi_{\lambda,g}^{\rho,\tau_0-\varepsilon\tau_1}(\alpha) &= - \int_{S_0/T} \pi^* \alpha \\
&= - \int_{\mathcal{M}_0/T_0} \text{Res}_\infty \left(\frac{\alpha}{\prod_{\nu \notin I} e_T(\mathcal{IND}(\bar{\partial}^\nu))} \right) \\
&= \int_{\mathcal{M}_0/T_0} \pi_0^* \alpha_0,
\end{aligned}$$

where

$$\alpha_0(\xi) = \frac{1}{2\pi i} \oint \frac{\alpha(\xi + ze_1)}{\prod_{\nu \notin I} \langle w_\nu, \xi + ze_1 \rangle^{d_\nu+1-g}} \exp \left(\sum_{\nu \notin I} \frac{\Omega_\nu}{\langle w_\nu, \xi + ze_1 \rangle} \right) dz$$

(by Lemma 4.10). This proves the theorem. $\square$

Proof of Lemma 4.10. Note first that $\mathbb{L}^\nu$ is the pullback under the projection $\pi_0 \times \text{id} : \mathcal{B}_I \times \Sigma \rightarrow \mathcal{A}/\mathcal{G}_0 \times \Sigma$ of the bundle

$$\mathcal{L}^\nu := \frac{\mathcal{A} \times L_\nu}{\mathcal{G}_0} \rightarrow \mathcal{A}/\mathcal{G}_0 \times \Sigma,$$

and $\mathcal{IND}(\bar{\partial}^\nu) \in K_T(\mathcal{B}_I)$ is the pullback under π_0 of the index bundle of the Cauchy-Riemann operators on $\mathcal{L}^\nu$. The torus T acts trivially on $\mathcal{A}/\mathcal{G}_0 \times \Sigma$ and by ρ_ν on the fibres of $\mathcal{L}^\nu$. Hence by Example 4.5,

$$e_T(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)) = \sum_{j \geq 0} \langle w_\nu, \xi \rangle^{d_\nu+1-g-j} c_j(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)). \quad (24)$$

Hence it remains to compute the *ordinary* Chern classes of the K -theory class $\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)$. The Atiyah–Singer index theorem for families asserts that

$$\text{ch}(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)) = \int_\Sigma \text{td}(T\Sigma) \text{ch}(\mathcal{L}^\nu) \in H^*(\mathcal{A}/\mathcal{G}).$$

(See [1, Theorem 4.3] and [2, Theorem 5.1].) Here ch and td denote the Chern character and the Todd class, respectively. The Todd class of $T\Sigma$ is given by

$$\text{td}(T\Sigma) = 1 + (1 - g)\sigma,$$

where $\sigma \in H^2(\Sigma; \mathbb{Z})$ denotes the positive generator. Thus our task at hand is to compute the Chern character of the line bundle $\mathcal{L}^\nu$. By Lemma 4.11 below, the first Chern class of $\mathcal{L}^\nu$ is given by

$$c_1(\mathcal{L}^\nu) = d_\nu \sigma - \sum_{i=1}^{2g} \sum_{j=1}^k w_{\nu j} \alpha_i \wedge \tau_{ij}.$$

From this we can compute $\text{ch}(\mathcal{L}^\nu)$. Note that $\alpha_i \wedge \alpha_{i'} = \pm \sigma$ whenever $i' = i \pm g$ and is equal to zero otherwise. Hence

$$\begin{aligned} \frac{1}{2} \left(\sum_{i=1}^{2g} \sum_{j=1}^k w_{\nu j} \alpha_i \wedge \tau_{ij} \right)^2 &= -\sigma \wedge \sum_{i=1}^g \sum_{j,j'=1}^k w_{\nu j} w_{\nu j'} \tau_{ij} \wedge \tau_{i+g,j'} \\ &=: -\sigma \wedge \Omega_\nu, \end{aligned}$$

and all higher powers vanish. It follows that

$$\text{ch}(\mathcal{L}^\nu) = 1 + d_\nu \sigma - \sum_{i=1}^{2g} \sum_{j=1}^k w_{\nu j} \alpha_i \wedge \tau_{ij} - \sigma \wedge \Omega_\nu.$$

Applying the index theorem for families we obtain

$$\begin{aligned} \text{ch}(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)) &= \int_\Sigma \text{td}(T\Sigma) \text{ch}(\mathcal{L}^\nu) \\ &= \int_\Sigma \left((d_\nu + 1 - g) \sigma - \sum_{i=1}^{2g} \sum_{j=1}^k w_{\nu j} \alpha_i \wedge \tau_{ij} - \sigma \wedge \Omega_\nu \right) \\ &= d_\nu + 1 - g - \Omega_\nu. \end{aligned}$$

The last formula implies, by a standard algebraic argument, that

$$c_1(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)) = -\Omega_\nu, \quad c_j(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)) = \frac{1}{j!} c_1(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu))^j.$$

Hence, by (24),

$$\begin{aligned} e_T(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu)) &= \sum_{j \geq 0} \langle w_\nu, \xi \rangle^{d_\nu + 1 - g - j} \frac{(-\Omega_\nu)^j}{j!} \\ &= \langle w_\nu, \xi \rangle^{d_\nu + 1 - g} \exp \left(-\frac{\Omega_\nu}{\langle w_\nu, \xi \rangle} \right). \end{aligned}$$

Since $e_T(\mathcal{IND}(\bar{\partial}^\nu)) = \pi_0^* e_T(\mathcal{IND}(\bar{\partial}^\nu, \mathcal{L}^\nu))$ the result follows. $\square$

Lemma 4.11. *The first Chern class of $\mathcal{L}^\nu$ is given by*

$$c_1(\mathcal{L}^\nu) = d_\nu \sigma - \sum_{i=1}^{2g} \sum_{j=1}^k w_{\nu j} \alpha_i \wedge \tau_{ij}.$$

Proof. Fix a reference connection $A_0 \in \mathcal{A}(P)$ and denote by $\mathcal{A}_0 \subset \mathcal{A}(P)$ the set of connections $A \in \mathcal{A}(P)$ that satisfy $F_A = \text{Vol}(\Sigma)^{-1} \lambda(P) \text{dvol}_\Sigma$ and $d^*(A - A_0) = 0$. The restricted gauge group $\mathcal{G}_{00} \subset \mathcal{G}_0(P)$ consists of all gauge

transformations $g : \Sigma \rightarrow T$ that satisfy $d^*(g^{-1}dg) = 0$ and $g(z_0) = \mathbb{1}$. Identify the quotient $\mathcal{A}_0/\mathcal{G}_{00}$ with the standard torus $\mathbb{T}^{2gk}$ via the map

$$\mathbb{R}^{2gk} \rightarrow \mathcal{A}_0 : t \mapsto A_t := A_0 + \sum_{i=1}^{2g} \sum_{j=1}^k t_{ij} \alpha_i e_j.$$

For i and j let $g_{ij} \in \mathcal{G}_{00}$ be the unique harmonic gauge transformation that satisfies $g_{ij}^{-1}dg_{ij} = \alpha_i e_j$ (and $g_{ij}(z_0) = \mathbb{1}$). Then the restriction of $\mathcal{L}^\nu$ to the submanifold $\mathcal{A}_0/\mathcal{G}_{00} \times \Sigma$ can be identified with the quotient $\mathbb{R}^{2gk} \times L_\nu/\mathbb{Z}^{2gk}$, where $m = \{m_{ij}\} \in \mathbb{Z}^{2gk}$ acts by

$$m^*(t, z, v) := (t + m, z, \prod_{ij} \rho_\nu(g_{ij}(z))^{-m_{ij}} v).$$

A section is a map $\mathbb{R}^{2gk} \times \Sigma \rightarrow L_\nu : (t, z) \mapsto s(t, z) = s_t(z) \in L_{\nu z}$ that satisfies

$$s(t + m, z) = \prod_{ij} \rho_\nu(g_{ij}(z))^{-m_{ij}} s(t, z), \quad m \in \mathbb{Z}^{2gk}.$$

A connection is given by the formula

$$d^\nabla s := d_{A_t} s_t + \sum_{i=1}^{2g} \sum_{j=1}^k \frac{\partial s_t}{\partial t_{ij}} dt_{ij}.$$

An easy computation shows that the curvature of this connection is the 2-form $F^\nabla \in \Omega^2(\mathbb{T}^{2gk} \times \Sigma, \sqrt{-1}\mathbb{R})$ given by

$$\begin{aligned} F^\nabla &= \dot{\rho}_\nu(F_{A_t}) - \sum_{i=1}^{2g} \sum_{j=1}^k \dot{\rho}_\nu \left(\frac{\partial A_t}{\partial t_{ij}} \right) \wedge dt_{ij} \\ &= -2\pi\sqrt{-1} \langle w_\nu, \lambda(P) \rangle \frac{\text{dvol}_\Sigma}{\text{Vol}(\Sigma)} + \sum_{i=1}^{2g} \sum_{j=1}^k 2\pi\sqrt{-1} \langle w_\nu, e_j \rangle \alpha_i \wedge dt_{ij}. \end{aligned}$$

Since the first Chern class of $\mathcal{L}^\nu$ is represented by the 2-form $\sqrt{-1}F^\nabla/2\pi$, the result follows. $\square$

5 Computation of the genus zero invariants

Let $\mathcal{I}$ denote the set of partitions

$$I_1 \cup \dots \cup I_k = \{1, \dots, n\}$$

that satisfy the (Dimension) condition in the introduction, i.e. for every $j \in \{1, \dots, k\}$, the subspace

$$E_j := \text{span}\{w_\nu \mid \nu \in I_1 \cup \dots \cup I_j\} \subset \mathfrak{t}^*$$

has dimension j and $w_\nu \notin E_j$ for every $\nu \in I_{j+1} \cup \dots \cup I_k$. It is now convenient to fix an orientation of $\mathfrak{t}$. For every $I = (I_1, \dots, I_k) \in \mathcal{I}$ we introduce the iterated residue $\phi_\lambda^I = \phi_\lambda^{\rho, I} : S^*(\mathfrak{t}^*) \rightarrow \mathbb{R}$ by

$$\phi_\lambda^I(\alpha) := \frac{1}{(2\pi i)^k} \oint \dots \oint \frac{\alpha(\sum z_j e_j)}{\prod_{\nu=1}^n \langle w_\nu, \sum z_j e_j \rangle^{d_\nu+1}} dz_k \dots dz_1, \quad (25)$$

where $d_\nu := \langle w_\nu, \lambda \rangle$. Here the lattice vectors $e_1, \dots, e_k \in \mathfrak{t}$ form an oriented integral basis of Λ such that the vectors $e_j, \dots, e_k$ are orthogonal to the span of the vectors w_ν for $\nu \in I_1 \cup \dots \cup I_{j-1}$ and $2 \leq j \leq k$. These requirements determine the e_j up to a change $e'_j = \pm e_j + \sum_{i>j} a_{ij} e_i$. The corresponding coordinates $\xi = \sum z_j e_j = \sum z'_j e'_j$ change by $z_i = \pm z'_i + \sum_{j<i} a_{ij} z'_j$. Since the e_i and the e'_i form oriented bases there is an even number of minus signs. It follows from these observations and Remark 3.2 that the integral (25) is independent of the choice of the e_i .

Lemma 5.1. *For every regular value τ of μ there exists a collection of integers $\{m_I\}_{I \in \mathcal{I}}$ such that*

$$\Phi_\lambda^{\rho, \tau}(\alpha) = \sum_{I \in \mathcal{I}} m_I \phi_\lambda^I(\alpha)$$

for every $\lambda \in \Lambda$ and every $\alpha \in S^*(\mathfrak{t}^*)$. Moreover, given a path γ connecting τ to $\mathfrak{t}^* \setminus \text{im } \mu$, the coefficients m_I can be chosen to satisfy the following condition: If $I = (I_1, \dots, I_k) \in \mathcal{I}$ is such that γ does not intersect the hyperplane spanned by the w_ν for $\nu \in I_1 \cup \dots \cup I_{k-1}$, then $m_I = 0$.

Proof. Theorem 1.1 and induction over the dimension of T . $\square$

Remark 5.2. Fix an oriented basis $e_1, \dots, e_k$ of $\mathfrak{t}$, let $H_\nu \subset \mathbb{C}^k$ be the hyperplane $\sum_{j=1}^k \langle w_\nu, e_j \rangle z_j = 0$, and denote $H := \bigcup_{\nu=1}^n H_\nu$. Then the right hand side of (25) can be interpreted as the integral of the k -form

$$\omega_{\lambda, \alpha} := \frac{\alpha(\sum z_j e_j)}{(2\pi i)^k \prod_{\nu=1}^n \langle w_\nu, \sum z_j e_j \rangle^{\langle w_\nu, \lambda \rangle + 1}} dz_1 \wedge \dots \wedge dz_k \in \Omega^k(\mathbb{C}^k \setminus H)$$

over a suitable homology class $\sigma_I \in H_k(\mathbb{C}^k \setminus H)$. Hence, by Lemma 5.1, there is a locally constant map $\tau \mapsto \sigma(\tau) := \sum_{I \in \mathcal{I}} m_I(\tau) \sigma_I$ which assigns to every regular value of μ a homology class $\sigma(\tau) \in H_k(\mathbb{C}^k \setminus H)$ such that the invariant $\Phi_\lambda^{\rho, \tau}(\alpha)$ is equal to the integral of $\omega_{\lambda, \alpha}$ over $\sigma(\tau)$ (for all λ and α). It is an interesting problem to study the map $\tau \mapsto \sigma(\tau)$ in more detail.

Lemma 5.3. *For $\nu = 1, \dots, n$ let ℓ_ν be integers and a_ν, b_ν be real numbers such that $a_\nu \neq 0$. If $\sum_{\nu=1}^n \ell_\nu = -1$ then*

$$\frac{1}{2\pi i} \oint \prod_{\nu=1}^n (a_\nu z + b_\nu)^{\ell_\nu} dz = \prod_{\nu=1}^n a_\nu^{\ell_\nu}.$$

If $\sum_{\nu=1}^n \ell_\nu < -1$ then the integral is zero.

Proof. In the variable $w := 1/z$ the integrand reads

$$\prod_{\nu} \left(\frac{a_{\nu}}{w} + b_{\nu} \right)^{\ell_{\nu}} \left(\frac{-dw}{w^2} \right) = - \frac{\prod_{\nu} (a_{\nu} + b_{\nu} w)^{\ell_{\nu}} dw}{w^{2+\sum \ell_{\nu}}}.$$

Since $a_{\nu} \neq 0$ the numerator is holomorphic near the origin. Hence the residue is zero whenever $2 + \sum_{\nu} \ell_{\nu} \leq 0$ and is equal to $\prod_{\nu} a_{\nu}^{\ell_{\nu}}$ whenever $2 + \sum_{\nu} \ell_{\nu} = 1$. $\square$

Lemma 5.4. *Let $I \in \mathcal{I}$ and $e_1, \dots, e_k$ be a positive basis of Λ such that $e_j, \dots, e_k$ are orthogonal to w_{ν} for $\nu \in I_1 \cup \dots \cup I_{j-1}$ (as in (25)). Let $\lambda \in \Lambda$ and ℓ be an n -tuple of nonnegative integers such that*

$$|\ell| = n - k + \sum_{\nu=1}^n d_{\nu}, \quad d_{\nu} := \langle w_{\nu}, \lambda \rangle. \quad (26)$$

If $I \in \mathcal{I}_{\lambda}(\ell)$ then

$$\phi_{\lambda}^I(w^{\ell}) = \prod_{j=1}^k \prod_{\nu \in I_j} \langle w_{\nu}, e_j \rangle^{\ell_{\nu} - d_{\nu} - 1}.$$

Otherwise $\phi_{\lambda}^I(w^{\ell}) = 0$.

Proof. The condition $I \in \mathcal{I}_{\lambda}(\ell)$ asserts that

$$\sum_{\nu \in I_j} (\ell_{\nu} - d_{\nu} - 1) = -1$$

for $j = 1, \dots, k$. Consider the integral over z_k . The coefficient $\langle w_{\nu}, e_k \rangle$ of z_k in the linear map $(z_1, \dots, z_k) \mapsto \langle w_{\nu}, \sum_j z_j e_j \rangle$ is nonzero iff $\nu \in I_k$. So, by Lemma 5.3,

$$\begin{aligned} \phi_{\lambda}^I(w^{\ell}) &= \left(\prod_{\nu \in I_k} \langle w_{\nu}, e_k \rangle^{\ell_{\nu} - d_{\nu} - 1} \right) \frac{1}{(2\pi i)^{k-1}} \\ &\quad \oint \cdots \oint \prod_{j=1}^{k-1} \prod_{\nu \in I_j} \langle w_{\nu}, \sum_{j=1}^{k-1} z_j e_j \rangle^{\ell_{\nu} - d_{\nu} - 1} dz_{k-1} \cdots dz_1 \end{aligned}$$

whenever $\sum_{\nu \in I_k} (\ell_{\nu} - d_{\nu} - 1) = -1$. If $\sum_{\nu \in I_k} (\ell_{\nu} - d_{\nu} - 1) < -1$ then the integral over z_k is zero. Hence it follows by induction that $\phi_{\lambda}^I(w^{\ell})$ has the required form whenever $I \in \mathcal{I}_{\lambda}(\ell)$. If $I \notin \mathcal{I}_{\lambda}(\ell)$ then it follows from (26) that $\sum_{\nu \in I_j} (\ell_{\nu} - d_{\nu} - 1) < -1$ for some j and hence $\phi_{\lambda}^I(w^{\ell}) = 0$. $\square$

Proof of Theorem 1.2. Assertion (i) follows from Lemmata 5.1 and 5.4.

We prove (ii). Let $\lambda \in \Lambda$, $d_{\nu} := \langle w_{\nu}, \lambda \rangle$, ℓ be an n -tuple of nonnegative integers, $J \subset \{1, \dots, n\}$ be an index set such that $\{w_{\nu} \mid \nu \in J\}$ is a basis of $\mathfrak{t}^*$, and assume that $\ell_{\nu} = d_{\nu}$ for $\nu \in J$ and $\ell_{\nu} = d_{\nu+1}$ for $\nu \notin J$. Then a partition I belongs to $\mathcal{I}_{\lambda}(\ell)$ if and only if $I \in \mathcal{I}$ and $I_j \cap J$ consists of a single element for

each j . This follows from the fact that $\sum_{\nu \in I_j} (\ell_\nu - d_\nu - 1) = -1$ and that each summand $\ell_\nu - d_\nu - 1$ equals 0 or -1 .

Assume $\tau \notin C(J) = \{\sum_{\nu \in J} \eta_\nu \mathbf{w}_\nu \mid \eta_\nu \geq 0\}$. We must prove that $\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = 0$. To see this, we examine the set $\mathcal{I}_\lambda(\ell)$. Since the set $\{\mathbf{w}_\nu \mid \nu \in J\}$ is linearly independent it follows that, for each ordering $J = \{\nu_1, \dots, \nu_k\}$, there exists a unique partition $I \in \mathcal{I}_\lambda(\ell)$ such that $\nu_j \in I_j$ for all j , and conversely, each partition $I \in \mathcal{I}_\lambda(\ell)$ determines an ordering of J . Moreover, for every such partition the hyperplane $W_I := \text{span}\{\mathbf{w}_\nu \mid \nu \notin I_k\}$ agrees with the hyperplane $W_{\nu_k} := \text{span}\{\mathbf{w}_\nu \mid \nu \in J \setminus \{\nu_k\}\}$. Hence the hyperplanes W_I for $I \in \mathcal{I}_\lambda(\ell)$ are precisely the supporting hyperplanes of $C(J)$. Since $\tau \notin C(J)$, there exists a straight line γ connecting τ to $\mathfrak{t}^* \setminus \text{im } \mu$ which misses the supporting hyperplanes (this is true because $C(J)$ is a cone over a simplex). Hence the coefficients m_I in Lemma 5.1 can be chosen such that $m_I = 0$ for every $I \in \mathcal{I}_\lambda(\ell)$. This implies that

$$\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = \sum_{I \in \mathcal{I}_\lambda(\ell)} m_I \phi_\lambda^I(\mathbf{w}^\ell) = 0.$$

The same argument shows that the invariant $\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell)$ for $\tau \in C(J)$ is independent of τ .

Assume $\tau \in C(J)$. Fix an ordering $J = \{\nu_1, \dots, \nu_k\}$ and let $I \in \mathcal{I}_\lambda(\ell)$ be the unique partition satisfying $\nu_j \in I_j$ for $j = 1, \dots, k$. Choose an integer basis $e_1, \dots, e_k$ of $\mathfrak{t}$ such that $\langle \mathbf{w}_\nu, e_j \rangle = 0$ for $\nu \in I_1 \cup \dots \cup I_{j-1}$ and $\langle \mathbf{w}_{\nu_j}, e_j \rangle > 0$. Let τ_0 be a positive linear combination of $\mathbf{w}_{\nu_1}, \dots, \mathbf{w}_{\nu_{k-1}}$. Since the invariant is independent of the choice of $\tau \in C(J)$, we may assume $\tau = \tau_0 + \varepsilon \tau_1$, where $\tau_1 := \mathbf{w}_{\nu_k}$. Since the invariant is zero outside of $C(J)$, we have

$$\Phi_\lambda^{\rho, \tau_0 - \varepsilon \tau_1}(\mathbf{w}^\ell) = 0.$$

Hence, by Theorem 1.1,

$$\Phi_\lambda^{\rho, \tau}(\mathbf{w}^\ell) = \frac{1}{\langle \mathbf{w}_{\nu_k}, e_k \rangle} \Phi_{\lambda_0}^{\rho_0, \tau_0} \left(\prod_{j=1}^{k-1} \prod_{\nu \in I_j} \mathbf{w}_\nu^{\ell_\nu} \right).$$

Now assertion (ii) follows by induction.

We prove (iii). Assume $|\ell| = n - k + \sum_{\nu=1}^n d_\nu$ (otherwise both invariants are zero). Since

$$\ell_\nu - d_\nu - 1 = \ell_\nu + d'_\nu - (d_\nu + d'_\nu) - 1,$$

for every ν we have

$$\mathcal{I}_\lambda(\ell) = \mathcal{I}_{\lambda+\lambda'}(\ell + d'), \quad \phi_\lambda^I(\mathbf{w}^\ell) = \phi_{\lambda+\lambda'}^I(\mathbf{w}^{\ell+d'})$$

for every $I \in \mathcal{I}$ (see Lemma 5.4). Hence (iii) follows from Lemma 5.1.

To prove (iv) and (v) we introduce the following notation. For every $\lambda \in \Lambda$ and every n -tuple ℓ of nonnegative integers define the number

$$\iota_\lambda(\ell) := \sum_{d_\nu \geq 0} \max\{\ell_\nu - d_\nu - 1, 0\} + \sum_{d_\nu \leq -1} \ell_\nu, \quad d_\nu := \langle \mathbf{w}_\nu, \lambda \rangle.$$

Recall that $J_\ell := \{\nu \mid \ell_\nu \leq d_\nu\}$. We prove (iv) and (v) in five steps.

Step 1. *Each w^ℓ can be expressed as a linear combination of classes $w^{\ell'}$ that satisfy $J_{\ell'} \subset J_\ell$ and either $\mathcal{I}_\lambda(\ell') = \emptyset$ or $\ell'_\nu \leq \max\{d_\nu + 1, 0\}$ for all ν .*

We prove Step 1 by induction over $\iota_\lambda(\ell)$. If $\iota_\lambda(\ell) = 0$ then $\ell'_\nu \leq \max\{d_\nu + 1, 0\}$ for all ν . Assume $\iota_\lambda(\ell) > 0$ and, by induction, that the claim has been established for every ℓ' that satisfies $\iota_\lambda(\ell') < \iota_\lambda(\ell)$. If $\mathcal{I}_\lambda(\ell) = \emptyset$ there is nothing to prove. Hence assume $\mathcal{I}_\lambda(\ell) \neq \emptyset$. Since $\iota_\lambda(\ell) > 0$ there is a ν_0 such that $\ell_{\nu_0} > 0$ and $\ell_{\nu_0} > d_{\nu_0} + 1$. Let $I = (I_1, \dots, I_k) \in \mathcal{I}_\lambda(\ell)$. Since $\sum_{\nu \in I_j} (\ell_\nu - d_\nu - 1) = -1$ for every j there are indices $\nu_j \in I_j$ for $j = 1, \dots, k$ such that $\ell_{\nu_j} \leq d_{\nu_j}$. By the (Dimension) condition, the vectors $w_{\nu_1}, \dots, w_{\nu_k}$ form a basis of $\mathfrak{t}^*$. Hence w_{ν_0} can be expressed as a linear combination of the vectors w_{ν_j} . Since $\ell_{\nu_0} > 0$ we can replace one of the factors w_{ν_0} in w^ℓ by this linear combination. This expresses w^ℓ as a linear combination of monomials of the form $w^{\ell'}$ with $\iota_\lambda(\ell') < \iota_\lambda(\ell)$ and $J_{\ell'} \subset J_\ell$. Hence the assertion for w^ℓ follows from the induction hypothesis.

Step 2. *If $\ell_\nu \leq d_\nu + 1$ for every ν and $\mathcal{I}_\lambda(\ell) \neq \emptyset$ then ℓ satisfies (ii).*

Let $I \in \mathcal{I}_\lambda(\ell)$. Then the formula $\sum_{\nu \in I_j} (\ell_\nu - d_\nu - 1) = -1$ shows that, for each j , there is precisely one index $\nu_j \in I_j$ such that $\ell_{\nu_j} = d_{\nu_j}$ and $\ell_\nu = d_\nu + 1$ for $\nu \in I_j \setminus \{\nu_j\}$. Since the vectors $w_{\nu_1}, \dots, w_{\nu_k}$ form a basis of $\mathfrak{t}^*$, it follows that ℓ satisfies (ii) with $J = \{\nu_1, \dots, \nu_k\}$.

Step 3. *We prove (iv).*

Assume $d_\nu \geq -1$ for every ν . Then, by Step 1, each $w^\ell \in S^{m_\lambda}(\mathfrak{t}^*)$ is a linear combination of classes $w^{\ell'}$ that satisfy either $\mathcal{I}_\lambda(\ell') = \emptyset$ or $\ell'_\nu \leq d_\nu + 1$ for all ν . Hence the assertion follows from Step 2.

Step 4. *(v) holds under the assumption $d_\nu \geq -1$ for all ν .*

We argue indirectly and assume that $\Phi_\lambda^{\rho, \tau}(w^\ell) \neq 0$. Then the linear combination in Step 1 must contain a term $w^{\ell'}$ that satisfies $J_{\ell'} \subset J_\ell$, and $\Phi_\lambda^{\rho, \tau}(w^{\ell'}) \neq 0$. The latter implies that $\mathcal{I}_\lambda(\ell') \neq \emptyset$ and so $\ell'_\nu \leq d_\nu + 1$ for all ν . Hence, by Step 2, ℓ' satisfies (ii) with $J = J_{\ell'} = \{\nu \mid \ell'_\nu \leq d_\nu\}$. Since $\Phi_\lambda^{\rho, \tau}(w^{\ell'}) \neq 0$, it follows from (ii) that $\tau \in C(J_{\ell'}) \subset C(J_\ell)$.

Step 5. *We prove (v).*

Suppose $\Phi_\lambda(w^\ell) \neq 0$. Choose λ' such that $d'_\nu := \langle w_\nu, \lambda' \rangle \geq \max\{0, -1 - d_\nu\}$ for all ν . Then, by (iii), we have $\Phi_{\lambda+\lambda'}(w^{\ell+d'}) = \Phi_\lambda(w^\ell) \neq 0$. Hence, by Step 4, $\tau \in C(J)$, where $J := \{\nu \mid \ell_\nu + d'_\nu \leq \langle w_\nu, \lambda + \lambda' \rangle\} = \{\nu \mid \ell_\nu \leq d_\nu\}$. This proves the theorem. $\square$

6 Quantum cohomology

Let τ be a regular value of μ . Throughout this section we assume that T acts freely on $\mu^{-1}(\tau)$. Equivalently, if $J \subset \{1, \dots, n\}$ is an index set consisting of k elements such that $\tau \in C(J)$, then τ belongs to the interior of $C(J)$ and the determinant of the tuple $\{w_\nu \mid \nu \in J\}$ is equal to plus or minus one. Under this assumption the symplectic quotient

$$\bar{M} := M // T(\tau) = \mu^{-1}(\tau) / T$$

is a Kähler manifold. We denote by $H^*(\bar{M})$, respectively $H_*(\bar{M})$, the quotients of the integral (co)homology groups by their torsion subgroups. By Kirwan's theorem, the homomorphism $H^*(BT) \rightarrow H^*(\bar{M})$ is surjective and the homomorphism $H_*(\bar{M}) \rightarrow H_*(BT)$ is injective. For every ν denote by $\bar{w}_\nu \in H^2(\bar{M})$ the image of the cohomology class $w_\nu \in \Lambda^* \cong H^2(BT)$ under the Kirwan homomorphism $H^2(BT) \rightarrow H^2(\bar{M})$. By Theorem C.4, the cohomology class $\bar{w}_\nu$ vanishes whenever $\tau \notin C(\{1, \dots, n\} \setminus \{\nu\})$. The remaining classes $\bar{w}_\nu$ generate $H^2(\bar{M})$. Hence the image of the homomorphism $H_2(\bar{M}) \rightarrow H_2(BT) \cong \Lambda$ is the subgroup

$$\Lambda(\tau) := \{\lambda \in \Lambda \mid \tau \notin C(\{1, \dots, n\} \setminus \{\nu\}) \implies \langle w_\nu, \lambda \rangle = 0\}.$$

Recall the definition of the inverse isomorphism $\Lambda(\tau) \rightarrow H_2(\bar{M}) : \lambda \mapsto \bar{\lambda}$ and the effective cone $\Lambda_{\text{eff}}(\tau) \subset \Lambda(\tau)$

$$\Lambda_{\text{eff}}(\tau) := \{\lambda \in \Lambda(\tau) \mid \langle \tau', \lambda \rangle \geq 0 \text{ for all } \tau' \in C(\tau)\},$$

where $C(\tau)$ denotes the chamber of τ . Note that

$$\langle \tau', \lambda \rangle > 0 \quad \text{for } \lambda \in \Lambda_{\text{eff}}(\tau) \setminus \{0\}, \quad \tau' \in C(\tau).$$

Denote by $\mathcal{D}_{\text{eff}}(\tau) \subset \mathbb{Z}^n$ the cone

$$\mathcal{D}_{\text{eff}}(\tau) := \{(\langle w_1, \lambda \rangle, \dots, \langle w_n, \lambda \rangle) \mid \lambda \in \Lambda_{\text{eff}}(\tau)\}.$$

Note that the map $\Lambda_{\text{eff}}(\tau) \rightarrow \mathcal{D}_{\text{eff}}(\tau)$ is a bijection. We denote the inverse by $\mathcal{D}_{\text{eff}}(\tau) \rightarrow \Lambda_{\text{eff}}(\tau) : d \mapsto \lambda_d$. We emphasize that $\mathcal{D}_{\text{eff}}(\tau)$ is not necessarily contained in the positive quadrant of $\mathbb{Z}^n$.

Let us now consider the vector

$$\tau := \sum_{\nu=1}^n w_\nu. \tag{27}$$

(We still assume that T acts freely on $\mu^{-1}(\tau)$.) Then $\bar{M}$ is a monotone symplectic manifold (see Lemma C.3). The genus zero Gromov-Witten invariants of $\bar{M}$ with fixed marked points in a homology class $\bar{\lambda} \in H_2(\bar{M})$ are denoted by

$$\text{GW}_{\bar{\lambda}}^{\bar{M}} : H^*(\bar{M}) \times \dots \times H^*(\bar{M}) \rightarrow \mathbb{Z}.$$

The number of arguments will in each case be clear from the context. For an n -tuple $\ell = (\ell_1, \dots, \ell_n)$ of nonnegative integers and a cohomology class $\bar{\alpha} \in H^*(\bar{M})$ we abbreviate

$$\text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{\ast \ell}, \bar{\alpha}) := \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}_1, \dots, \bar{w}_1, \dots, \bar{w}_n, \dots, \bar{w}_n, \bar{\alpha}),$$

where each argument $\bar{w}_\nu$ occurs ℓ_ν times. Since the Gromov-Witten invariants are invariant under symplectic deformation, we have

$$\lambda \in \Lambda(\tau) \setminus \Lambda_{\text{eff}}(\tau) \implies \text{GW}_{\bar{\lambda}}^{\bar{M}} \equiv 0.$$

Lemma 6.1. *Let $\tau := \sum_{\nu=1}^n w_\nu$, suppose that T acts freely on $\mu^{-1}(\tau)$, and assume that the minimal Chern number N of M is greater than one. Then for every $d \in \mathcal{D}_{\text{eff}}(\tau)$, every $\lambda \in \Lambda(\tau)$, and every $\bar{\alpha} \in H^*(M)$, we have*

$$\text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{*d^+}, \bar{\alpha}) = \text{GW}_{\bar{\lambda}-\bar{\lambda}_d}^{\bar{M}}(\bar{w}^{*d^-}, \bar{\alpha}), \quad (28)$$

where the n -tuples d^+ and d^- are defined by

$$d_\nu^+ := \begin{cases} d_\nu, & \text{if } d_\nu > 0, \\ 0, & \text{if } d_\nu \leq 0, \end{cases} \quad d_\nu^- := \begin{cases} -d_\nu, & \text{if } d_\nu < 0, \\ 0, & \text{if } d_\nu \geq 0. \end{cases}$$

Proof. Let $\bar{\alpha} := \bar{w}^\ell$ for an n -tuple ℓ of nonnegative integers satisfying

$$|\ell| = n - k + \langle \tau, \lambda \rangle - |d^+|.$$

By Theorem 1.2 (iii), with λ replaced by $\lambda - \lambda_d$, $\lambda' = \lambda_d$, and ℓ replaced by $\ell + d^-$, we have

$$\Phi_{\lambda}^{\rho, \tau}(w^{d^++\ell}) = \Phi_{\lambda-\lambda_d}^{\rho, \tau}(w^{d^-+\ell}). \quad (29)$$

Hence it follows from [8, Theorem A] and the fact that $N > 1$ (see equation (3) in the introduction) that

$$\text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{*(d^++\ell)}) = \text{GW}_{\bar{\lambda}-\bar{\lambda}_d}^{\bar{M}}(\bar{w}^{*(d^-+\ell)}). \quad (30)$$

Now the gluing theorem for the Gromov–Witten invariants with fixed marked points (see [11]) asserts that

$$\begin{aligned} \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{*d^+}, \bar{w}^\ell) &= \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{*(d^++\ell)}) \\ &\quad - \sum_i \sum_{\lambda' \neq 0} \text{GW}_{\bar{\lambda}-\bar{\lambda}'}^{\bar{M}}(\bar{w}^{*d^+}, \bar{e}_i) \text{GW}_{\bar{\lambda}'}^{\bar{M}}(\bar{e}_i^*, \bar{w}^{*\ell}), \end{aligned}$$

where the second sum is over all lattice vectors $\lambda' \in \Lambda_{\text{eff}}(\tau) \setminus \{0\}$. Hence, by (30),

$$\begin{aligned} &\text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{*d^+}, \bar{w}^\ell) - \text{GW}_{\bar{\lambda}-\bar{\lambda}_d}^{\bar{M}}(\bar{w}^{*d^-}, \bar{w}^\ell) \\ &= \sum_i \sum_{\lambda' \neq 0} \text{GW}_{\bar{\lambda}'}^{\bar{M}}(\bar{e}_i^*, \bar{w}^{*\ell}) \left(\text{GW}_{\bar{\lambda}-\bar{\lambda}'}^{\bar{M}}(\bar{w}^{*d^+}, \bar{e}_i) - \text{GW}_{\bar{\lambda}-\bar{\lambda}'}^{\bar{M}}(\bar{w}^{*d^-}, \bar{e}_i) \right). \end{aligned} \quad (31)$$

Note that in each summand on the right we have

$$\begin{aligned} \frac{1}{2} \deg(e_i) &= n - k + \langle \tau, \lambda - \lambda' \rangle - |d^+| \\ &< n - k + \langle \tau, \lambda \rangle - |d^+| \\ &= |\ell|. \end{aligned}$$

Hence the assertion follows from (31) by induction over $|\ell|$. $\square$

Remark. Let $\tau, \bar{M}$ be as in Lemma 6.1 and $\lambda \in \Lambda_{\text{eff}}(\tau)$ such that $\langle w_\nu, \lambda \rangle \geq 0$ for every ν . Then it follows from Lemma 6.1 with $d_\nu := \langle w_\nu, \lambda \rangle$ and $\bar{\alpha} := \text{PD}(\text{pt})$ that $\text{GW}_{\bar{\lambda}}^{\bar{M}} \neq 0$. Hence the homology class $\bar{\lambda} \in H_2(\bar{M})$ can be represented by a holomorphic stable map of genus zero.

As in the introduction, let $\mathcal{R}$ be any graded commutative algebra (over the reals) with unit which is equipped with a homomorphism

$$\Lambda_{\text{eff}}(\tau) \rightarrow \mathcal{R} : \lambda \mapsto q^\lambda$$

from the additive semigroup $\Lambda_{\text{eff}}(\tau)$ to the multiplicative semigroup $\mathcal{R}$ such that

$$\deg(q^\lambda) = 2\langle \tau, \lambda \rangle.$$

The most important example is the ring

$$\mathcal{R} = \mathbb{R}[q_1, \dots, q_k, q_1^{-1}, \dots, q_k^{-1}]$$

of polynomials with real coefficients in the variables q_j and q_j^{-1} . To obtain the homomorphism choose a basis $e_1, \dots, e_k$ of Λ , define the grading by $\deg(q_j) = \sum_{\nu=1}^n \langle w_\nu, e_j \rangle$, and the map $\lambda \mapsto q^\lambda$ by

$$q^\lambda := \prod_{j=1}^k q_j^{\lambda_j}, \quad \lambda = \sum_{j=1}^k \lambda_j e_j.$$

With a more careful choice of the basis one can take $\mathcal{R} = \mathbb{R}[q_1, \dots, q_k]$. Other possibilities are the polynomial ring $\mathcal{R} = \mathbb{R}[q]$ in one variable, the ring of polynomials in q and q^{-1} , or the ring of Laurent series in q . In these cases one can choose q to have degree two and define $q^\lambda := q^{\langle \tau, \lambda \rangle}$. The simplest example is $\mathcal{R} = \mathbb{R}$ with the constant map $\lambda \mapsto q^\lambda := 1$, but then the grading has to be reduced modulo $2N$, where N is the minimal Chern number.

Given a graded algebra $\mathcal{R}$ as above define the quantum cohomology ring $\text{QH}^*(\bar{M}; \mathcal{R})$ as the tensor product

$$\text{QH}^*(\bar{M}; \mathcal{R}) := H^*(\bar{M}; \mathbb{R}) \otimes \mathcal{R}$$

(of vector spaces over the reals). Thus an element of $\text{QH}^m(\bar{M}; \mathcal{R})$ is a finite sum $\bar{\alpha} = \sum_{r \in \mathcal{R}} \bar{\alpha}_r r$ such that $\deg(\bar{\alpha}_r) + \deg(r) = m$ for all r . The ring structure is defined by

$$\bar{\alpha}' * \bar{\alpha}'' := \sum_i \sum_{\lambda \in \Lambda_{\text{eff}}(\tau)} \sum_{r', r''} \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{\alpha}'_i, \bar{\alpha}''_i, \bar{e}_i) \bar{e}_i^* r' r'' q^\lambda,$$

where the $\bar{e}_i$ form a basis of $H^*(\bar{M})$ and the $\bar{e}_i^*$ denote the dual basis with respect to the cup product pairing (see [11]).

Corollary 6.2. *Let $\tau := \sum_{\nu=1}^n w_\nu$, suppose that T acts freely on $\mu^{-1}(\tau)$, and assume that the minimal Chern number N of $\bar{M}$ is greater than one. Then*

$$\bar{w}^* d^+ = \bar{w}^* d^- q^{\lambda_d}$$

for every $d \in \mathcal{D}_{\text{eff}}(\tau)$.

Proof. By the gluing theorem for the Gromov–Witten invariants (see [11]), we have

$$\begin{aligned}\bar{w}^{*d^+} &= \sum_{i,\lambda} \text{GW}_{\bar{\lambda}}^{\bar{M}}(\bar{w}^{*d^+}, \bar{e}_i) \bar{e}_i^* q^\lambda \\ &= \sum_{i,\lambda} \text{GW}_{\bar{\lambda}-\bar{\lambda}_d}^{\bar{M}}(\bar{w}^{*d^-}, \bar{e}_i) \bar{e}_i^* q^\lambda \\ &= \bar{w}^{*d^-} q^{\lambda_d}.\end{aligned}$$

The second equality follows from Lemma 6.1. $\square$

Proof of Theorem 1.3. We prove that the homomorphism (4) is surjective. Note that there is an obvious inclusion $H^*(\bar{M}) \rightarrow \text{QH}^*(\bar{M}; \mathcal{R}) : \bar{\alpha} \mapsto \bar{\alpha}1$, where 1 denotes the unit in $\mathcal{R}$. Throughout we identify $H^*(\bar{M})$ with its image in $\text{QH}^*(\bar{M}; \mathcal{R})$ under this homomorphism. Since (4) is a homomorphism of $\mathcal{R}$ -modules, it suffices to prove that every class in $H^*(\bar{M})$ belongs to the image of (4). We prove this by induction over the degree. If $\bar{\alpha} \in H^0(\bar{M})$ then $\bar{\alpha}$ obviously belongs to the image of (4). Hence let $\deg(\bar{\alpha}) = 2\ell > 0$ and assume, by induction, that every class in $H^*(\bar{M})$ of degree less than 2ℓ belongs to the image of (4). By Kirwan’s theorem, the class $\bar{\alpha}$ is a linear combination of classes of the form $\bar{w}_{\nu_1} \cdots \bar{w}_{\nu_\ell}$. Let $p(u_1, \dots, u_n)$ be the same linear combination of the polynomials $u_{\nu_1} \cdots u_{\nu_\ell}$. Then the image of $p(u)$ under the homomorphism differs from $\bar{\alpha}$ by a class of the form

$$\bar{\beta} = \sum_{\lambda \neq 0} \bar{\beta}_\lambda q^\lambda, \quad \deg(\bar{\beta}_\lambda) = 2\ell - 2\langle \tau, \lambda \rangle < 2\ell.$$

Here the sum is over all $\lambda \in \Lambda_{\text{eff}}(\tau)$ that satisfy $\langle \tau, \lambda \rangle > 0$. Hence, by the induction hypothesis, every $\bar{\beta}_\lambda$ in this sum belongs to the image of (4), and so does the class $\bar{\beta}_\lambda q^\lambda$. Hence $\bar{\beta}$ belongs to the image of (4), and so does $\bar{\alpha}$.

Let $\mathcal{J}_0 \subset \mathcal{R}[u_1, \dots, u_n]$ be the kernel of (4). Then the linear polynomial $\sum_\nu \eta_\nu u_\nu$ belongs to $\mathcal{J}_0$ whenever $\sum_\nu \eta_\nu w_\nu = 0$. Moreover, by Corollary 6.2, the polynomial $u^{d^+} - q^\lambda u^{d^-}$ belongs to $\mathcal{J}_0$ whenever $\lambda \in \Lambda_{\text{eff}}(\tau)$ and $d_\nu^\pm = \max\{\pm \langle w_\nu, \lambda \rangle, 0\}$. Hence $\mathcal{J} \subset \mathcal{J}_0$.

We prove that $\mathcal{J}_0 \subset \mathcal{J}$. Define the classes $\bar{\alpha}_{\ell,\lambda} \in H^*(\bar{M})$, for n -tuples ℓ of nonnegative integers and lattice vectors $\lambda \in \Lambda_{\text{eff}}(\tau)$ with $0 < \langle \tau, \lambda \rangle \leq |\ell|$, by

$$\bar{w}^{*\ell} =: \bar{w}^\ell + \sum_{\lambda \in \Lambda_{\text{eff}}(\tau) \setminus \{0\}} \bar{\alpha}_{\ell,\lambda} q^\lambda, \quad \deg(\bar{\alpha}_{\ell,\lambda}) = 2|\ell| - 2\langle \tau, \lambda \rangle.$$

For $N \in \mathbb{Z}$ denote by $\mathcal{J}_0(N)$ the set of polynomials $p \in \mathcal{J}_0$ of the form

$$p(u_1, \dots, u_n) = \sum_{|\ell| \leq N} r_\ell u^\ell, \tag{32}$$

where the sum is over all n -tuples $\ell = (\ell_1, \dots, \ell_n)$ of nonnegative integers satisfying $|\ell| \leq N$. We prove by induction on N that $\mathcal{J}_0(N) \subset \mathcal{J}$. For $N < 0$

this is obvious because $\mathcal{J}_0(N) = \{0\}$. Let $N \geq 0$ and assume by induction that $\mathcal{J}_0(N-1) \subset \mathcal{J}$. Let $p \in \mathcal{J}_0(N)$ be a polynomial of the form (32). Since $p \in \mathcal{J}_0$ we have

$$0 = \sum_{\ell} r_{\ell} \bar{w}^{*\ell} = \sum_{\ell} \bar{w}^{\ell} r_{\ell} + \sum_{\ell} \sum_{\lambda} \bar{\alpha}_{\ell, \lambda} r_{\ell} q^{\lambda}.$$

This identity splits up into

$$0 = \sum_{|\ell|=j} r_{\ell} \bar{w}^{\ell} + \sum_{|\ell|>j} \sum_{\substack{\lambda \\ \langle \tau, \lambda \rangle = |\ell| - j}} \bar{\alpha}_{\ell, \lambda} r_{\ell} q^{\lambda}, \quad j = 0, \dots, N.$$

Since $r_{\ell} = 0$ for $|\ell| > N$, we have

$$\sum_{|\ell|=N} r_{\ell} \bar{w}^{\ell} = 0.$$

Choose a basis $\rho_1, \dots, \rho_m$ of the vector space $\text{span}\{r_{\ell} \mid |\ell| = N\} \subset \mathcal{R}$ and express each r_{ℓ} in this basis, i.e.

$$r_{\ell} = \sum_{i=1}^m a_{\ell i} \rho_i, \quad a_{\ell i} \in \mathbb{R}, \quad |\ell| = N.$$

Then

$$\sum_{|\ell|=N} a_{\ell i} \bar{w}^{\ell} = 0, \quad i = 1, \dots, m.$$

This means that the polynomials

$$p_{i0}(u_1, \dots, u_n) := \sum_{|\ell|=N} a_{\ell i} u^{\ell}, \quad i = 1, \dots, m,$$

belong to the kernel $\mathcal{I} \subset \mathbb{R}[u_1, \dots, u_n]$ of the homomorphism (36) in Theorem C.4. Hence they can be expressed in the form

$$p_{i0} = \sum_j p_{ij0} f_j,$$

where $f_j \in \mathbb{R}[u_1, \dots, u_n]$, and the p_{ij0} are taken from the set of generators of $\mathcal{I}$ in Theorem C.5. Thus each p_{ij0} satisfies one of the following conditions.

- (a) $p_{ij0}(u) = \sum_{\nu} \eta_{\nu} u_{\nu}$, where $\sum_{\nu} \eta_{\nu} w_{\nu} = 0$.
- (b) $p_{ij0}(u) = u_{\nu}$, where $\Delta_{\{\nu\}} = \emptyset$.
- (c) $p_{ij0}(u) = u^{d^+}$, where $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$.

In cases (a) and (b) define $p_{ij} := p_{ij0} \in \mathcal{J}$. In the case (c) it follows from the definition of $\mathcal{J}$ that there is a generator $p_{ij} \in \mathcal{J}$ of the form

$$p_{ij}(u) = u^{d^+} - q^{\lambda} u^{d^-}, \quad p_{ij0}(u) = u^{d^+}, \quad d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}.$$

Define $\hat{p} \in \mathcal{J}$ by

$$\hat{p}(u) := \sum_{i=1}^m \sum_j \rho_i p_{ij}(u) f_j(u).$$

Since $\mathcal{J} \subset \mathcal{J}_0$ we have $p - \hat{p} \in \mathcal{J}_0$. Since

$$\sum_{i=1}^m \sum_j \rho_i p_{ij0}(u) f_j(u) = \sum_{i=1}^m \rho_i p_{i0}(u) = \sum_{i=1}^m \sum_{|\ell|=N} \rho_i a_{\ell i} u^\ell = \sum_{|\ell|=N} r_\ell u^\ell,$$

the leading terms cancel in $p - \hat{p}$ and hence $p - \hat{p} \in \mathcal{J}_0(N-1) \subset \mathcal{J}$. Hence $p \in \mathcal{J}$. This completes the induction and the proof of the theorem. $\square$

Example 6.3. This example shows that in the definition of the ideal $\mathcal{J}$ it may not suffice to consider vectors $\lambda \in \Lambda_{\text{eff}}(\tau)$ such that the integers $d_\nu := \langle w_\nu, \lambda \rangle$ are all nonnegative. Suppose the 2-torus $T = \mathbb{T}^2$ acts on $\mathbb{C}^5$ with weight vectors

$$w_1 = (1, 0), \quad w_2 = (1, 1), \quad w_3 = w_4 = w_5 = (0, 1).$$

The symplectic quotient $\bar{M}$ at the parameter $\tau := w_1 + \dots + w_5 = (2, 4)$ is a smooth monotone toric 3-fold with minimal Chern number $N = 2$. The effective cone is given by

$$\Lambda_{\text{eff}}(\tau) = \{(\lambda_1, \lambda_2) \in \mathbb{Z}^2 \mid \lambda_2 \geq 0, \lambda_1 + \lambda_2 \geq 0\}.$$

It is the convex cone spanned by the vectors $e := (1, 0)$ with $d = (1, 1, 0, 0, 0)$ and $e' = (-1, 1)$ with $d' = (-1, 0, 1, 1, 1)$. For the quantum cohomology let us choose the polynomial ring $\mathcal{R} := \mathbb{R}[q_1, q_2]$, graded by $\deg(q_1) = \deg(q_2) = 4$, and the homomorphism $q^\lambda := q_1^{\lambda_1 + \lambda_2} q_2^{\lambda_2}$. Thus q_1, q_2 correspond to the generators e, e' of $\Lambda_{\text{eff}}(\tau)$. Then the ideal $\mathcal{J} \subset \mathbb{R}[u_1, \dots, u_5, q_1, q_2]$ is generated by the relations

$$u_3 = u_4 = u_5 = u_2 - u_1, \quad u_1 u_2 = q_1, \quad u_3 u_4 u_5 = u_1 q_2. \quad (33)$$

If one considers only vectors $\lambda \in \Lambda_{\text{eff}}(\tau)$ with nonnegative degrees $d_\nu := \langle w_\nu, \lambda \rangle$ then one has to replace the last relation in (33) by $u_2 u_3 u_4 u_5 = q_1 q_2$ and obtains a strictly smaller ideal.

A Equivariant trivialization

Proposition A.1. *Let G be a compact Lie group and $E \rightarrow X$ be a G -equivariant complex vector bundle over compact smooth manifold X . Then there exists a G -equivariant complex vector bundle $F \rightarrow X$ and a complex G -representation W such that $E \oplus F$ is equivariantly isomorphic to $X \times W$.*

Lemma A.2. *Let G be a compact Lie group, $H \subset G$ be a normal subgroup, and V be a complex H -representation. Then there exists a complex G -representation W and an injective H -equivariant homomorphism $\phi : V \rightarrow W$.*

Proof. Consider the infinite dimensional vector space

$$\mathcal{W} := \{f : G \rightarrow V \mid f(hg) = hf(g) \ \forall h \in H, \ \forall g \in G\}.$$

This space carries an action of G by

$$(g'f)(g) := f(gg')$$

and the evaluation map $\mathcal{W} \rightarrow V : f \mapsto f(1)$ is H -equivariant and surjective. To prove surjectivity let $v \in V$ be given and let $f : G \rightarrow V$ be any smooth extension of the map $H \rightarrow V : h \mapsto hv$. By averaging the maps $g \mapsto h^{-1}f(hg)$ over $h \in H$ we can ensure that the extension is H -equivariant. By Peter–Weyl’s theorem, there exists a finite dimensional G -invariant subspace $W \subset \mathcal{W}$ such that the restriction of the homomorphism $f \mapsto f(1)$ to W is still surjective. By [11, Remark A.4.2], the surjection $W \rightarrow V$ has an H -equivariant right inverse. $\square$

Proof of Proposition A.1. Let $x \in X$, $H \subset G$ be the isotropy subgroup of x , and $V := E_x$. By the local slice theorem, the restriction of E to a suitable neighbourhood of the G -orbit of x is equivariantly isomorphic to the bundle

$$\frac{G \times U \times V}{H} \rightarrow G \times_H U,$$

where U is a neighbourhood of zero in the horizontal tangent space at x (i.e. in the orthogonal complement of $T_x Gx$ with respect to some G -invariant metric). Let $\phi : V \rightarrow W$ be as in Lemma A.2. Then the map

$$G \times U \times V \rightarrow G \times U \times W : (g, u, v) \mapsto (g, u, g\phi(v))$$

descends to a G -equivariant injective bundle homomorphism from $(G \times U \times V)/H$ to $(G \times_H U) \times W$ (where G acts diagonally). This construction gives rise to a G -invariant open cover $\{U_\alpha\}_\alpha$ of X and a collection of G -equivariant injective bundle homomorphisms $\phi_\alpha : E|_{U_\alpha} \rightarrow U_\alpha \times W_\alpha$. Let $\rho_\alpha : X \rightarrow [0, 1]$ be a G -invariant partition of unity subordinate to the cover $\{U_\alpha\}_\alpha$ and denote $W := \bigoplus_\alpha W_\alpha$. Then the homomorphism $E \rightarrow X \times W : (x, e) \mapsto (x, \{\rho_\alpha(x)\phi_\alpha(x)e\}_\alpha)$ is the required G -equivariant embedding. $\square$

B Convex polytopes

In this section we recall some well-known facts about convex polytopes (see e.g. [7]). Let Δ be a compact convex polytope in the dual space V^* of a finite dimensional vector space V . We denote elements of V by v, w and elements of V^* by ξ, η . Define the *support function* $\phi : V \rightarrow \mathbb{R}$ of Δ by

$$\phi(v) := \inf_{\xi \in \Delta} \langle \xi, v \rangle.$$

The following properties of ϕ are obvious from the definition.

(P1) $\phi(tv) = t\phi(v)$ for $t \geq 0$.

(P1) ϕ is concave, i.e. $\phi(v + w) \geq \phi(v) + \phi(w)$.

(P3) Δ can be recovered from ϕ as the intersection of half spaces

$$\Delta = \bigcap_{v \in V} \{\xi \in V^* \mid \langle \xi, v \rangle \geq \phi(v)\}.$$

Let F be a face of Δ . Pick an interior point p of F and define the *dual cone to F* by

$$\check{F} := \{v \in V \mid \langle \xi - p, v \rangle \geq 0 \text{ for all } \xi \in \Delta\}.$$

If q is another interior point of F and $\xi \in \Delta$ then $q + t(\xi - p) \in \Delta$ for $t > 0$ sufficiently small. Hence $\langle \xi - p, v \rangle \geq 0$ iff $\langle q + t(\xi - p) - q, v \rangle \geq 0$. This shows that the definition of $\check{F}$ does not depend on the point p . Moreover, the condition $\langle \xi - p, v \rangle \geq 0$ for all $\xi \in \Delta$ can be rewritten as $\phi(v) \geq \langle p, v \rangle$, or equivalently $\phi(v) = \langle p, v \rangle$ since $p \in \Delta$. So $\check{F}$ can be written in the equivalent forms

$$\begin{aligned} \check{F} &= \{v \in V \mid \langle \xi - p, v \rangle \geq 0 \text{ for all } \xi \in \Delta, p \in F\} \\ &= \{v \in V \mid \langle p, v \rangle = \phi(v) \text{ for all } p \in F\}. \end{aligned}$$

The following properties are obvious from these descriptions of $\check{F}$.

(F1) $\check{F}$ is a convex polyhedral cone.

(F2) The restriction of ϕ to $\check{F}$ is the linear function $\phi(v) = \langle p, v \rangle$ for any $p \in F$.

(F3) $\check{F}$ is perpendicular to F and $\dim \check{F} = \text{codim} F$.

(F4) If $H_1, \dots, H_\ell \subset V^*$ are the supporting hyperplanes for Δ meeting at F , then $\check{F}$ is the cone generated by inward pointing normal vectors $v_1, \dots, v_\ell$ to the hyperplanes.

(F5) If G is a subface of F then $\check{F}$ is a subcone of $\check{G}$.

(F6) The union of the cones $\check{p}$ dual to vertices p of Δ is the whole space V .

The collection Σ of the cones $\check{F}$ dual to nonempty faces of Δ is called the *fan dual to Δ* (see [3] for the general definition of a fan).

C The cohomology of symplectic quotients

Let T be a k -dimensional torus and $\rho = (\rho_1, \dots, \rho_n) : T \rightarrow \mathbb{T}^n := (S^1)^n$ be a diagonal homomorphism with

$$\rho_\nu(\exp(\xi)) = e^{-2\pi i \langle w_\nu, \xi \rangle}$$

for $\xi \in \mathfrak{t} := \text{Lie}(T)$. Here the w_ν are elements of the dual lattice $\Lambda^* \subset \mathfrak{t}^*$ as in the introduction. We identify the Lie algebra of $\mathbb{T}^n$ with $\mathbb{R}^n$ via the map $\eta \mapsto i\eta/2\pi$

so that the integer lattice corresponds to $\mathbb{Z}^n \subset \mathbb{R}^n$. In this identification the linearization of ρ is the map $\dot{\rho} : \mathfrak{t} \rightarrow \mathbb{R}^n$ given by

$$\dot{\rho}(\xi) = (\langle w_1, \xi \rangle, \dots, \langle w_n, \xi \rangle).$$

Consider the quotient torus

$$\bar{T} := \mathbb{T}^n / \rho(T).$$

Its Lie algebra is the quotient space

$$\bar{\mathfrak{t}} := \mathbb{R}^n / \dot{\rho}(\mathfrak{t})$$

and the dual space of $\bar{\mathfrak{t}}$ can be identified with the subspace

$$\bar{\mathfrak{t}}^* := \left\{ \eta \in (\mathbb{R}^n)^* \mid \sum_{\nu=1}^n \eta_\nu w_\nu = 0 \right\}. \quad (34)$$

The canonical action of $\mathbb{T}^n$ on $\mathbb{C}^n$ induces an action of T with moment map $\mu : \mathbb{C}^n \rightarrow \mathfrak{t}^*$ given by (1). We assume throughout that μ is proper and that the action is effective (i.e. the weight vectors w_ν span $\mathfrak{t}^*$). Let $\tau \in \mathfrak{t}^*$ be a regular value of μ . Then the torus $\bar{T}$ acts on the symplectic quotient

$$\bar{M} := \mathbb{C}^n // T(\tau) = \mu^{-1}(\tau) / T.$$

A moment map $\bar{\mu} : \bar{M} \rightarrow \bar{\mathfrak{t}}^*$ for this action is given by the formula

$$\bar{\mu}(x) := \begin{pmatrix} \pi|x_1|^2 + \zeta_1 \\ \vdots \\ \pi|x_n|^2 + \zeta_n \end{pmatrix},$$

where $\zeta = (\zeta_1, \dots, \zeta_n) \in (\mathbb{R}^n)^*$ is chosen such that

$$\sum_{\nu=1}^n \zeta_\nu w_\nu = -\tau.$$

The image of $\bar{\mu}$ is the convex polyhedron

$$\Delta := \bar{\mu}(\bar{M}) = \left\{ \eta \in (\mathbb{R}^n)^* \mid \sum_{\nu=1}^n \eta_\nu w_\nu = 0, \eta_\nu \geq \zeta_\nu \right\}. \quad (35)$$

Each subset $I \subset \{1, \dots, n\}$ determines a (possibly empty) face

$$\Delta_I := \{ \eta \in \Delta \mid \eta_\nu = \zeta_\nu \text{ for } \nu \in I \}.$$

Recall that $C(I)$ denotes the cone spanned by the vectors w_ν , $\nu \in I$. The next lemma shows that if τ is a regular value of μ , then the intersection of any j codimension-1 faces of Δ is either empty or has codimension j .

Lemma C.1. Assume that τ is a regular value of μ and let $I \subset \{1, \dots, n\}$.

(i) The set Δ_I is either empty or has codimension $|I|$.

(ii) $\Delta_I = \emptyset \iff \tau \notin C(\{1, \dots, n\} \setminus I)$.

Proof. We prove (i). Assume $\Delta_I \neq \emptyset$ and let $J := \{1, \dots, n\} \setminus I$. Then, by the definition of $\bar{\mu}$, there is a $y \in \mathbb{C}^J$ such that

$$\mu(y) = \pi \sum_{\nu \in J} |y_\nu|^2 w_\nu = \tau.$$

Since τ is a regular value of μ , there exist indices $\nu_1, \dots, \nu_k \in J$ such that the vectors $w_{\nu_1}, \dots, w_{\nu_k}$ are linearly independent and $y_{\nu_j} \neq 0$ for every j . We claim that there is a vector $x \in \mathbb{C}^J$ such that

$$\mu(x) = \tau, \quad x_\nu \neq 0 \quad \text{for all } \nu \in J.$$

To see this choose x_ν for $\nu \in J \setminus \{\nu_1, \dots, \nu_k\}$ such that $|x_\nu|^2 = |y_\nu|^2 + \varepsilon$ and choose x_{ν_j} such that

$$\sum_{j=1}^k (|x_{\nu_j}|^2 - |y_{\nu_j}|^2) w_{\nu_j} + \varepsilon \sum_{\nu \in J \setminus \{\nu_1, \dots, \nu_k\}} w_\nu = 0.$$

Then $\mu(x) = \mu(y) = \tau$ and, for $\varepsilon > 0$ sufficiently small, we get $x_\nu \neq 0$ for all $\nu \in J$.

The differential $d\bar{\mu}(x) : T_x(\bar{M} \cap \mathbb{C}^J) \rightarrow \bar{\mathfrak{t}}^*$ is given by

$$d\bar{\mu}(x)v = (2\pi \langle x_\nu, v_\nu \rangle)_{\nu \in J},$$

where $v \in \mathbb{C}^J$ satisfies

$$d\mu(x)v = 2\pi \sum_{\nu \in J} \langle x_\nu, v_\nu \rangle w_\nu = 0.$$

Since $x_\nu \neq 0$ for $\nu \in J$, this shows that the image of $T_x(\bar{M} \cap \mathbb{C}^J)$ under $d\bar{\mu}(x)$ equals $\{\eta \in (\mathbb{R}^J)^* \mid \sum_{\nu \in J} \eta_\nu w_\nu = 0\}$. This space, and therefore Δ_I , has dimension $|J| - k = n - k - |I|$.

We prove (ii). If $\Delta_I \neq \emptyset$ there exists an $\eta \in (\mathbb{R}^n)^*$ such that $\sum_{\nu=1}^n \eta_\nu w_\nu = 0$, $\eta_\nu \geq \zeta_\nu$ for all ν , and $\eta_\nu = \zeta_\nu$ for $\nu \in I$. Hence

$$\tau = \sum_{\nu=1}^n -\zeta_\nu w_\nu = \sum_{\nu \notin I} (\eta_\nu - \zeta_\nu) w_\nu \in C(\{1, \dots, n\} \setminus I).$$

The converse follows by reversing the argument. $\square$

Standing assumption. In the remainder of this appendix we assume that T acts freely on $\mu^{-1}(\tau)$.

Denote by $\bar{w}_\nu \in H^2(\bar{M}; \mathbb{R})$ the image of w_ν under the homomorphism $\Lambda^* \cong H^2(\text{BT}; \mathbb{Z}) \rightarrow H^*(\bar{M}; \mathbb{R})$.

Lemma C.2. *For every $J \subset \{1, \dots, n\}$ the following holds.*

(i) *If $\tau \notin C(J)$ then $\prod_{\nu \notin J} \bar{w}_\nu = 0$.*

(ii) *If $\tau \in C(J)$ and $|J| = k$ then $\prod_{\nu \notin J} \bar{w}_\nu = \text{PD}(\text{pt})$.*

Proof. $\bar{w}_\nu$ is the first Chern class of the line bundle $\bar{L}_\nu := \mu^{-1}(\tau) \times_{\rho_\nu} \mathbb{C}$. Hence the zero set of the holomorphic section $\bar{M} \rightarrow \bar{L}_\nu : [x] \mapsto [x, x_\nu]$ is Poincaré dual to $\bar{w}_\nu$. Denote this zero set by $\bar{W}_\nu := \{[x] \in \bar{M} \mid x_\nu = 0\}$. This is a (possibly empty) complex submanifold of $\bar{M}$ of complex codimension one. Moreover,

$$\tau \notin C(J) \implies \bigcap_{\nu \notin J} \bar{W}_\nu = \emptyset$$

This proves (i). If $\tau \in C(J)$ and $|J| = k$, then the submanifolds $\bar{W}_\nu$ for $\nu \notin J$ intersect transversally in a single point. This proves (ii). $\square$

Lemma C.3. (i) *The Chern classes of $T\bar{M}$ are given by*

$$c_j(T\bar{M}) = \sum_{\nu_1 < \dots < \nu_j} \bar{w}_{\nu_1} \cdots \bar{w}_{\nu_j}, \quad j = 1, \dots, n - k.$$

(ii) *The cohomology class of the symplectic form $\bar{\omega} \in \Omega^2(\bar{M})$ is*

$$[\bar{\omega}] = \bar{\tau}.$$

Proof. We prove (i). Consider the Whitney sum $\mu^{-1}(\tau) \times \mathbb{C}^n = E \oplus F$, where the complex vector bundles $E \rightarrow \mu^{-1}(\tau)$ and $F \rightarrow \mu^{-1}(\tau)$ are defined by

$$E_x := \{v \in \mathbb{C}^n \mid d\mu(x)v = d\mu(x)iv = 0\},$$

$$F_x := \{v \in \mathbb{C}^n \mid \exists \xi, \eta \in \mathfrak{t} \forall \nu : v_\nu = (\langle w_\nu, \xi \rangle + i \langle w_\nu, \eta \rangle) x_\nu\}.$$

Then the bundle F admits a T -equivariant complex trivialization and the quotient bundle $E/T \rightarrow \mu^{-1}(\tau)/T$ is isomorphic to the tangent bundle of $\bar{M}$. Hence

$$c_j(T\bar{M}) = c_j(\mu^{-1}(\tau) \times_T \mathbb{C}^n) = \sum_{\nu_1 < \dots < \nu_j} \bar{w}_{\nu_1} \cdots \bar{w}_{\nu_j}.$$

We prove (ii). Denote

$$\lambda_0 := \frac{1}{2i} \sum_{\nu=1}^n (\bar{x}_\nu dx_\nu - x_\nu d\bar{x}_\nu) \in \Omega^1(\mathbb{C}^n), \quad \omega_0 := d\lambda_0,$$

and let $\mu_0 : \mathbb{C}^n \rightarrow \mathbb{R}^n$ be the moment map given by

$$\mu_0(x) := \pi(|x_1|^2, \dots, |x_n|^2).$$

Since $d_{\mathbb{T}^n} \lambda_0 = \omega_0 - \mu_0$, the equivariant cohomology class $[\omega_0 - \mu_0] \in H_{\mathbb{T}^n}^2(\mathbb{C}^n)$ is trivial. Pulling back under the homomorphism $H_{\mathbb{T}^n}^2(\mathbb{C}^n) \rightarrow H_T^2(\mathbb{C}^n)$ induced by ρ yields $0 = [\omega_0 - \mu] \in H_T^2(\mathbb{C}^n)$. Restriction to $\mu^{-1}(\tau)$ yields $0 = [\iota^* \omega_0 - \iota^* \mu] \in H_T^2(\mu^{-1}(\tau))$, where $\iota : \mu^{-1}(\tau) \rightarrow \mathbb{C}^n$ is the inclusion. Now the result follows by passing to the quotient. $\square$

Theorem C.4 ([10]). *The ring homomorphism*

$$\mathbb{R}[u_1, \dots, u_n] \rightarrow H^*(\bar{M}; \mathbb{R}) : p(u_1, \dots, u_n) \mapsto p(\bar{w}_1, \dots, \bar{w}_n) \quad (36)$$

induces an isomorphism

$$H^*(\bar{M}; \mathbb{R}) \cong \mathbb{R}[u_1, \dots, u_n]/\mathcal{I},$$

where the ideal $\mathcal{I} \subset \mathbb{R}[u_1, \dots, u_n]$ is generated by the relations

$$\sum_{\nu=1}^n \eta_\nu w_\nu = 0 \quad \implies \quad \sum_{\nu=1}^n \eta_\nu u_\nu = 0, \quad (37)$$

$$I \subset \{1, \dots, n\}, \quad \Delta_I = \emptyset \quad \implies \quad \prod_{\nu \in I} u_\nu = 0. \quad (38)$$

Remark. Theorem C.4 continues to hold with coefficients in $\mathbb{Z}$. That the homomorphism (36) is surjective follows from Kirwan's theorem, and that the ideal $\mathcal{I}$ is contained in the kernel of (36) is an easy consequence of Lemma C.2. The nontrivial part of the proof is to show that the kernel is contained in $\mathcal{I}$.

Theorem C.5 ([4]). *Assume $\mu^{-1}(\tau) \neq \emptyset$ and T acts freely on $\mu^{-1}(\tau)$. Then the ideal $\mathcal{I}$ is generated by the linear relations (37), the linear monomials u_ν for $\Delta_{\{\nu\}} = \emptyset$, and the monomials u^d for $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$.*

For the sake of completeness, we present a somewhat more elaborated version of the proof given in [4]. We need some preparation. Denote by $\bar{\Lambda} \subset \bar{\mathfrak{t}}$ and $\bar{\Lambda}^* \subset \bar{\mathfrak{t}}^*$ the integer lattices. Thus $\bar{\Lambda}$ is the image of $\mathbb{Z}^n$ under the projection $\mathbb{R}^n \rightarrow \bar{\mathfrak{t}}$ and $\bar{\Lambda}^* = \bar{\mathfrak{t}}^* \cap (\mathbb{Z}^n)^*$. For $\nu = 1, \dots, n$ let $\bar{e}_\nu \in \bar{\Lambda}$ be the image of the basis vector $e_\nu = (0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{Z}^n$ under the projection $\mathbb{R}^n \rightarrow \bar{\mathfrak{t}}$.

Lemma C.6. *Suppose that T acts freely on $\mu^{-1}(\tau)$.*

- (i) *Let $J \subset \{1, \dots, n\}$ satisfy $|J| = n - k$ and $\Delta_J \neq \emptyset$. Then the vectors $\{\bar{e}_j \mid j \in J\}$ form an integer basis of $\bar{\Lambda}$.*
- (ii) *Let $d_\nu \in \mathbb{Z}$ satisfy $\sum_{\nu=1}^n d_\nu \bar{e}_\nu = 0$. Then there exists a vector $\lambda \in \Lambda$ such that $d_\nu = \langle w_\nu, \lambda \rangle$ for every ν .*

Proof. We prove (i). Assume $|J| = n - k$ and $\Delta_J \neq \emptyset$. Since T acts freely on $\mu^{-1}(\tau)$, and $\tau \in C(\{1, \dots, n\} \setminus J)$, the vectors $\{w_\nu \mid \nu \notin J\}$ form an integer basis of Λ^* . Hence, for every $v \in \mathbb{Z}^n$, there exists a unique vector $\lambda \in \Lambda$ such that $v_\nu = \langle w_\nu, \lambda \rangle$ for $\nu \notin J$. This implies that the image $\bar{v} \in \bar{\mathfrak{t}}$ of v under the projection $\mathbb{R}^n \rightarrow \bar{\mathfrak{t}}$ satisfies

$$\bar{v} = \sum_{j \in J} (v_j - \langle w_j, \lambda \rangle) \bar{e}_j.$$

Hence the vectors $\{\bar{e}_j \mid j \in J\}$ span the integer lattice $\bar{\Lambda}$ as claimed.

We prove (ii). By definition of the projection $\mathbb{R}^n \rightarrow \bar{\mathfrak{t}}$, there exists a vector $\xi \in \mathfrak{t}$ such that $d_\nu = \langle w_\nu, \xi \rangle$ for every ν . Now let $J \subset \{1, \dots, n\}$ be any index

set such that $|J| = n - k$ and $\Delta_J \neq \emptyset$. Then the argument in the proof of (i) shows that there exists a lattice vector $\lambda \in \Lambda$ such that $d_\nu = \langle w_\nu, \lambda \rangle$ for $\nu \notin J$. Hence $\langle w_\nu, \xi - \lambda \rangle = 0$ for $\nu \notin J$. Since the vectors $\{w_\nu \mid \nu \notin J\}$ form a basis of $\mathfrak{t}^*$ we deduce that $\xi = \lambda$ and hence $d_\nu = \langle w_\nu, \lambda \rangle$ for every ν . $\square$

Proof of Theorem C.5. Let $\mathcal{I}_0 \subset \mathbb{R}[u_1, \dots, u_n]$ be the ideal generated by the linear polynomials $\sum_{\nu=1}^n \eta_\nu u_\nu$, where $\sum_{\nu=1}^n \eta_\nu w_\nu = 0$, the monomials u_ν , where $\Delta_{\{\nu\}} = \emptyset$, and the monomials u^{d^+} for $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$.

We prove that $\mathcal{I}_0 \subset \mathcal{I}$. We must show that $u^{d^+} \in \mathcal{I}$ for every $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$. We prove a stronger statement: *If $\lambda \in \Lambda$ satisfies $\langle \tau, \lambda \rangle > 0$ and $d_\nu := \langle w_\nu, \lambda \rangle$ then $u^{d^+} \in \mathcal{I}$.* To see this, consider the set $I := \{\nu \mid d_\nu > 0\}$. We claim that $\Delta_I = \emptyset$. Otherwise, by Lemma C.1, there would exist numbers $\eta_\nu \geq 0$ such that $\tau = \sum_{\nu \notin I} \eta_\nu w_\nu$. But then

$$0 < \langle \tau, \lambda \rangle = \sum_{\nu \notin I} \eta_\nu d_\nu \leq 0,$$

a contradiction. Since $\Delta_I = \emptyset$, the monomial $\prod_{\nu \in I} u_\nu$ belongs to the ideal $\mathcal{I}$. But u^{d^+} is a multiple of $\prod_{\nu \in I} u_\nu$ and hence also belongs to $\mathcal{I}$.

We prove that $\mathcal{I} \subset \mathcal{I}_0$. Consider the moment polytope $\Delta \subset \bar{\mathfrak{t}}^*$ defined by (35). The faces of Δ are subsets of the form Δ_I for $I \subset \{1, \dots, n\}$ such that $\tau \in C(\{1, \dots, n\} \setminus I)$. The vectors $\{\bar{e}_i \mid i \in I\}$ are the inward pointing normal vectors to the supporting hyperplanes of Δ meeting at the face Δ_I . Hence, by property (F4) of the dual cones (see Appendix B), the dual cone of Δ_I is given by

$$\check{\Delta}_I = \left\{ \sum_{i \in I} c_i \bar{e}_i \mid c_i \geq 0 \right\}.$$

By Lemma C.1, the codimension of the face Δ_I equals $|I|$. In particular, the vertices of Δ are subsets Δ_J where $|J| = n - k$ and $\Delta_J \neq \emptyset$.

Now let $I \subset \{1, \dots, n\}$ such that $\Delta_I = \emptyset$. We must prove that the monomial $\prod_{\nu \in I} u_\nu$ belongs to $\mathcal{I}_0$. Shrinking the set I , if necessary, we may assume without loss of generality that $\Delta_{I'} \neq \emptyset$ for every proper subset $I' \subsetneq I$. Since $\mu(\tau) \neq \emptyset$ we have $I \neq \emptyset$. If $|I| = 1$ then the polynomial $\prod_{\nu \in I} u_\nu$ belongs to $\mathcal{I}_0$ by assumption. Hence assume $|I| \geq 2$. Then

$$\nu \in I \quad \implies \quad \Delta_{\{\nu\}} \neq \emptyset. \quad (39)$$

We shall prove that there exists a vector $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$ such that $d_\nu = 1$ for $\nu \in I$ and $d_\nu \leq 0$ for $\nu \notin I$. To see this, consider the vector $\sum_{i \in I} \bar{e}_i \in \bar{\Lambda}$. Since the union of the cones dual to vertices is the whole space $\bar{\mathfrak{t}}$, it follows that there exists an index set $J \subset \{1, \dots, n\}$ such that $|J| = n - k$, $\Delta_J \neq \emptyset$, and $\sum_{i \in I} \bar{e}_i \in \check{\Delta}_J$. Hence there exists nonnegative real numbers c_j such that

$$\sum_{i \in I} \bar{e}_i = \sum_{j \in J} c_j \bar{e}_j.$$

By Lemma C.6, the set $\{\bar{e}_j \mid j \in J\}$ is an integer basis of $\bar{\Lambda}$. Hence the c_j are actually integers and, after shrinking J , we may assume that $c_j > 0$ for all $j \in J$. Define $d \in \mathbb{Z}^n$ by

$$d_\nu := \begin{cases} 1, & \text{if } \nu \in I \setminus J, \\ -c_\nu, & \text{if } \nu \in J \setminus I, \\ 1 - c_\nu, & \text{if } \nu \in I \cap J, \\ 0, & \text{if } \nu \notin I \cup J. \end{cases}$$

Then $\sum_{\nu=1}^n d_\nu \bar{e}_\nu = 0$ and hence, by Lemma C.6, there exists a lattice vector $\lambda \in \Lambda$ such that $d_\nu = \langle w_\nu, \lambda \rangle$ for $\nu = 1, \dots, n$. By (39), we have $\Delta_{\{\nu\}} \neq \emptyset$ for every $\nu \in I \cup J$. Hence $d_\nu = 0$ whenever $\Delta_{\{\nu\}} = \emptyset$, and this implies $\lambda \in \Lambda(\tau)$.

We prove that $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$. Let $\phi : \mathfrak{t} \rightarrow \mathbb{R}$ be the support function of Δ as in Appendix B. By property (P2), we have

$$\sum_{i \in I} \phi(\bar{e}_i) \leq \phi\left(\sum_{i \in I} \bar{e}_i\right) = \phi\left(\sum_{j \in J} c_j \bar{e}_j\right) = \sum_{j \in J} c_j \phi(\bar{e}_j).$$

Here the last equation follows from property (F2) and the fact that the set $\{\bar{e}_j \mid j \in J\}$ spans the cone $\check{\Delta}_J$. Now, by (35) and the definition of ϕ , we have $\phi(\bar{e}_\nu) \geq \zeta_\nu$, with equality if and only if the face $\Delta_{\{\nu\}}$ is nonempty. Moreover, $d_\nu = 0$ whenever $\Delta_{\{\nu\}} = \emptyset$. This implies

$$0 \geq \sum_{\nu=1}^n d_\nu \phi(\bar{e}_\nu) = \sum_{\nu=1}^n d_\nu \zeta_\nu = \sum_{\nu=1}^n \langle w_\nu, \lambda \rangle \zeta_\nu = -\langle \tau, \lambda \rangle.$$

If we replace τ by another vector τ' in the same chamber, the fan Σ remains the same, so the above argument yields the same vector $\lambda \in \Lambda$. This shows that $\langle \tau', \lambda \rangle \geq 0$ for every τ' in the chamber of τ . So $\lambda \in \Lambda_{\text{eff}}(\tau)$ and $d \in \mathcal{D}_{\text{eff}}(\tau)$. Since $\Delta_I = \emptyset$ and $\Delta_J \neq \emptyset$, we have $I \neq J$ and hence $d \neq 0$.

We prove that $I \cap J = \emptyset$. Otherwise let $\nu_0 \in I \cap J$ and $I' := I \setminus \{\nu_0\}$. Then $d_\nu \leq 0$ for $\nu \notin I'$. Hence the argument in the proof of $\mathcal{I}_0 \subset \mathcal{I}$ shows that $\Delta_{I'} = \emptyset$. But this contradicts the minimality assumption on I . Hence $I \cap J = \emptyset$ as claimed. It follows that the vector d satisfies $d_\nu = 1$ for $\nu \in I$ and $d_\nu \leq 0$ for $\nu \notin I$. Since $d \in \mathcal{D}_{\text{eff}}(\tau) \setminus \{0\}$ we deduce that

$$\prod_{\nu \in I} u_\nu = u^{d^+} \in \mathcal{I}_0.$$

This shows that $\mathcal{I} \subset \mathcal{I}_0$ and hence $\mathcal{I} = \mathcal{I}_0$. □

References

- [1] M.F. Atiyah, I. Singer, The index of elliptic operators: III, *Annals of Mathematics* **87** (1968), 546–604.

- [2] M.F. Atiyah, I. Singer, The index of elliptic operators: IV, *Annals of Mathematics* **93** (1971), 119–138.
- [3] M. Audin, *The Topology of Torus Actions on Symplectic Manifolds*, Progress in Mathematics **93**, Birkhäuser, Basel 1991.
- [4] V. Batyrev, Quantum cohomology rings of toric manifolds, *Journées de Géométrie Algébrique d’Orsay* (Orsay, 1992), *Astisque* **218** (1993), 9–34.
- [5] K. Cieliebak, A.R. Gaio, I. Mundet i Rivera, D.A. Salamon, The symplectic vortex equations and invariants of Hamiltonian group actions. To appear in *J. Symp. Geom.*
- [6] K. Cieliebak, I. Mundet i Rivera, D.A. Salamon, Equivariant moduli problems and the Euler class. To appear in *Topology*.
- [7] T. Oda, *Convex Bodies and Algebraic Geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge Band 14, Springer, Berlin 1985.
- [8] A.R. Gaio, D.A. Salamon, Gromov–Witten invariants of symplectic quotients and adiabatic limits, Preprint ETH–Zürich, June 2001.
- [9] A.B. Givental, A mirror theorem for toric complete intersections, in *Topological Field Theory, Primitive Forms, and Related Topics*, Kyoto, 1996, pp. 141–175, Progr. Math. **160**, Birkhäuser, Boston 1998.
- [10] V. Guillemin, S. Sternberg, *Supersymmetry and Equivariant de Rham theory*, Springer, Berlin 1999.
- [11] D. McDuff, D. Salamon, *J-holomorphic Curves and Quantum Cohomology*, Third Edition, to appear 2003.
- [12] Z. Qin, Y. Ruan, Quantum cohomology of projective bundles over $\mathbb{P}^n$, *Trans. Amer. Math. Soc.* **350** (1998), 3615–3638.
- [13] H. Spielberg, The Gromov-Witten invariants of symplectic toric manifolds, Preprint, math.AG/006156 (2000).
- [14] H. Spielberg, Counting generic genus-0 curves on Hirzebruch surfaces, *Proc. Amer. Math. Soc.* **130** (2002), 1257–1264.